

D-A089 930

NAVAL POSTGRADUATE SCHOOL MONTEREY CA  
RELATIONSHIPS BETWEEN SEA SURFACE TEMPERATURE AND NUTRIENTS IN --ETC(U)  
MAR 80 J W CONRAD

F/6 17/5


UNCLASSIFIED


NL

1 of 2

2/28

1/1/77





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 089930

NAVAL POSTGRADUATE SCHOOL  
Monterey, California



DTIC  
ELECTE  
OCT 6 1980  
A

THESIS

RELATIONSHIPS BETWEEN SEA SURFACE TEMPERATURE  
AND NUTRIENTS IN SATELLITE DETECTED OCEANIC  
FRONTS

by

John Woeppel Conrad

March 1980

Thesis Advisor:

Eugene D. Traganza

Approved for public release; distribution unlimited

DDC FILE COPY

80 10 3 081

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A089	930
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
Relationships Between Sea Surface Temperature and Nutrients in Satellite Detected Oceanic Fronts.		Master's Thesis March 1980
6. AUTHOR(s)		7. PERFORMING ORG. REPORT NUMBER
John Woepfel/Conrad		
8. PERFORMING ORGANIZATION NAME AND ADDRESS		9. CONTRACT OR GRANT NUMBER(s)
Naval Postgraduate School Monterey, California 93940		(12) 1156
10. CONTROLLING OFFICE NAME AND ADDRESS		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Naval Postgraduate School Monterey, California 93940		
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. REPORT DATE
Naval Postgraduate School Monterey, California 93940		March 1980
		14. NUMBER OF PAGES
		114
		15. SECURITY CLASS. (of this report)
		Unclassified
		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Nutrients, Nitrate, Phosphate, satellite infrared imagery, thermal fronts, chemical fronts, fronts, Nitrogen, Phosphorus, sea surface temperatures.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Satellite IR images of the California coast off Point Sur reveal recurrent surface features which appear to be "thermal discontinuities" associated with aperiodic upwelling events. Some of these have associated "chemical fronts" and increased biological activity. Satellite IR imagery was used to locate "discontinuities" and with <u>in situ</u> monitoring the development of three features were studied. Interrelationships between sea surface temperature		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE/When Data Entered

20. (cont'd)

nutrients and microplanktonic biomass were investigated. Nutrient ratios, satellite imagery, wind stress data and correlations between nutrients and temperature were used to develop an estimate of "age" within a simplified upwelling "life cycle" model.

The features range in scale from tens to hundreds of kilometers. Two upwelling features exhibited very strong correlations between nutrient and temperature but a third feature had considerable nutrient variability. This suggests a considerable impact from the dynamic and biological processes. The technique of coupling satellite imagery and in situ monitoring was found to be a feasible method to provide real time inferences of the nutrient structure associated with an upwelled thermal feature.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE/When Data Entered

Approved for public release; distribution unlimited

Relationships Between Sea Surface Temperature  
and Nutrients in Satellite Detected  
Oceanic Fronts

by

John Woeppel Conrad  
Lieutenant Commander, United States Navy  
B.S., United States Naval Academy, 1969

Submitted in partial fulfillment of the  
requirements for the degree of

MASTER OF SCIENCE IN OCEANOGRAPHY

from the

NAVAL POSTGRADUATE SCHOOL  
March 1980

Author

*John Woeppel Conrad*

Approved by:

*Eugene D. Izaguirre*

Thesis Advisor

*Stevens P. Tucker*

Second Reader

*Glenn D. Doney*

for Chairman, Department of Oceanography

*William M. Tolles*

Dean of Science and Engineering

## ABSTRACT

Satellite IR images of the California coast off Point Sur reveal recurrent surface features which appear to be "thermal discontinuities" associated with aperiodic upwelling events. Some of these have associated "chemical fronts" and increased biological activity. Satellite IR imagery was used to locate "discontinuities" and with in situ monitoring the development of three features were studied. Interrelationships between sea surface temperature, nutrients and microplanktonic biomass were investigated. Nutrient ratios, satellite imagery, wind stress data and correlations between nutrients and temperature were used to develop an estimate of "age" within a simplified upwelling "life cycle" model.

The features range in scale from tens to hundreds of kilometers. Two upwelling features exhibited very strong correlations between nutrient and temperature but a third feature had considerable nutrient variability. This suggests a considerable impact from the dynamic and biological processes. The technique of coupling satellite imagery and in situ monitoring was found to be a feasible method to provide real time inferences of the nutrient structure associated with an upwelled thermal feature.

## TABLE OF CONTENTS

I.	INTRODUCTION-----	11
II.	THEORY-----	13
	A. UPWELLING-----	13
	B. SATELLITE IMAGERY-----	14
	C. NUTRIENTS (NITROGEN AND PHOSPHORUS)-----	15
III.	METHODS-----	17
	A. NUTRIENTS-----	17
	B. SATELLITE IMAGERY-----	17
	C. COMPUTATIONS-----	18
	D. CHLOROPHYLL-----	19
	E. ATP-----	19
	F. TEMPERATURE-----	19
	G. WIND-----	20
IV.	RESULTS-----	21
	A. 30 APRIL CRUISE-----	21
	B. 13 JUNE CRUISE-----	32
	C. 7 AUGUST CRUISE-----	33
V.	DISCUSSION-----	54
VI.	CONCLUSIONS-----	62
APPENDIX A.	Listing of Cruise Data: Time, Latitude, Longitude, Elapsed Distance, ATP, Nitrate, Phosphate, Chlorophyll, Nutrient Ratio, Temperature-----	64
APPENDIX B:	Wind Data: Wind Stress, Ekman Transport, Up- Welling Index, Vertical Velocity-----	102
BIBLIOGRAPHY-----		108
INITIAL DISTRIBUTION LIST-----		112



# LIST OF PHOTOGRAPHIC PLATES

Plate 1.	TIROS-N Satellite IR Image for 19 April 1979----	22
Plate 2.	TIROS-N Satellite IR Image for 29 April 1979----	23
Plate 3.	TIROS-N Satellite IR Image for 13 June 1979-----	34
Plate 4.	TIROS-N Satellite IR Image for 30 July 1979-----	42
Plate 5.	TIROS-N Satellite IR Image for 5 August 1979----	43

## LIST OF FIGURES

Fig. 1.	Track of the 30 April Cruise and Outline of Upwelling Feature-----	24
Fig. 2.	Nitrate, Phosphate and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 30 April Cruise-----	25
Fig. 3.	ATP, Chlorophyll <u>a</u> , and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 30 April Cruise-----	26
Fig. 4.	Nutrient Ratio and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 30 April Cruise-----	27
Fig. 5.	Nitrate Versus Phosphate for the 30 April Cruise-	28
Fig. 6.	Nitrate Versus Temperature for the 30 April Cruise-----	29
Fig. 7.	Phosphate Versus Temperature for the 30 April Cruise-----	30
Fig. 8.	Vertical Temperature Sections Along the Track of the 30 April Cruise-----	31
Fig. 9.	Track of the 13 June Cruise and Outline of the Upwelling Feature-----	35
Fig. 10.	Nitrate, Phosphate, Nutrient Ratio, ATP, Chlorophyll <u>a</u> and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 13 June Cruise-----	36
Fig. 11.	Nitrate Versus Phosphate for the 13 June Cruise--	37
Fig. 12.	Nitrate Versus Temperature for the 13 June Cruise-----	38
Fig. 13.	Phosphate Versus Temperature for the 13 June Cruise-----	39
Fig. 14.	Track of the 7 August Cruise and Outline of the Upwelling Feature-----	44

Fig. 15.	Nitrate, Phosphate and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise-----	45
Fig. 16.	ATP, Chlorophyll <u>a</u> , and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise-----	46
Fig. 17.	Nutrient Ratio and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise-----	47
Fig. 18.	Nitrate Versus Phosphate for the 7 August Cruise-	48
Fig. 19.	Nitrate Versus Temperature for the 7 August Cruise-----	49
Fig. 20.	Phosphate Versus Temperature for the 7 August Cruise-----	50
Fig. 21.	Vertical Temperature Sections Along the Track of the 7 August Cruise-----	51

## LIST OF TABLES

Table I.	Summary of Regression Analyses-----	52
Table II.	Data Ranges-----	53

## ACKNOWLEDGMENTS

This thesis is a result of ongoing research in chemical oceanography at the Naval Postgraduate School which is supported by the Office of Naval Research, Code 482, NSTL, Bay St. Louis, Mississippi. I thank the sponsors and key individuals whose assistance made this task possible, including: Dr. Eugene D. Traganza, my thesis advisor and principal investigator; Dr. Christopher N. K. Mooers, Dr. Stevens P. Tucker, Dr. Eugene C. Haderlie, Ms. Andrea McDonald of the Naval Postgraduate School, Monterey, California; Captain W. W. Reynolds and the crew of the R/V ACANIA; Mr. L. Breaker of the National Environmental Satellite Service at Redwood City, California; and Mr. A. Bakun of the National Marine Fisheries Service, Pacific Environmental Group, Monterey, California. Lastly, I thank my family for their forbearance, patience and support.

## I. INTRODUCTION

In this thesis satellite thermal imagery and in situ automated biochemical analyses were combined to study inter-relationships of sea surface temperatures, nitrate, phosphate and biological activity in oceanic fronts.

An oceanic front is a region of transition between two oceanic regimes with different characteristics (Cheney, 1976). These characteristics can be temperature, salinity, chemical or biological quantities.

In situ automated biochemical and thermal analyses of surface waters were previously accomplished by others, particularly Kelly, et al. (1975). However, they were not linked to satellite infrared imagery. Due to the marginal quality of satellite-derived results, oceanographers have largely either ignored them or have not been convinced that they could be integrated usefully with classical oceanographic observations (Legeckis, 1978). More sophisticated satellites and sensors, such as the TIROS (Television Infra-red Observation Satellites) series and the Advanced Very High Resolution Radiometer (AVHRR) have made it possible to make useful inferences in areas of the ocean that are classically associated with sea surface temperature anomalies. The region of interest in this study is off the central California coast, where "oceanic fronts" and "eddies" appear to form in response to wind driven pulses of cold upwelled

coastal waters. The existence of "chemical fronts" in association with these features was postulated in 1978 by Traganza (1978) and demonstrated in 1978 by Traganza et al. (1980).

Continued study of these frontal systems may be particularly relevant to the interests of the Navy in view of their potential effect on the propagation of sound energy and the capability of sensors to distinguish significant sound signals from the background noises in the ocean. Sound velocity changes across thermal fronts, coupled with changes in nutrient concentrations that lead to increased biological activity, may degrade both active and passive sonar performance through changes in sound propagation loss, increased biological reverberation levels and background noise levels.

## II. THEORY

### A. UPWELLING

Coastal upwelling is caused by the divergence of surface water away from the coast. Cold water rises from subsurface layers to replace this water. When the pycnocline surfaces, relatively nutrient-rich water (which has been trapped below the pycnocline due to density differences) will upwell and enrich the surface waters. In the California coastal area this water is generally derived from a depth no greater than 200 meters offshore (Fairbridge, 1966; Sverdrup, et al., 1942). In the Northern Hemisphere coastal upwelling can be caused by the northerly wind stress prevailing in spring and summer along the western continental coasts. These prevailing winds may vary seasonally off the California coast as the North Pacific subtropical high moves north during the spring and summer, then south during the autumn and winter (Bakun, 1973). Upwelling can also be caused by currents impinging on land masses (Fairbridge, 1966) or winds blowing directly away from shore.

The circulation patterns vary with the width and form of the continental shelf and slope, the wind speed, duration, fetch and latitude. The characteristics of upwelled water may vary depending on the depth from which the water came, the properties of the source water, the characteristic vertical velocity, and the residence time of the upwelled



water in the euphotic zone (SCOR Working Group, 1974). The proximity of the heads of underwater canyons enhances the intensity of the upwelling, (SCOR Working Group, 1974; Codispoti, 1977; Treguer, 1977). Therefore, upwelling can vary from region to region and season to season (Codispoti et al., 1979; Barton et al., 1977).

#### B. SATELLITE IMAGERY

Infrared (IR) images of the sea surface temperatures were received from the TIROS-N series satellites using an Advanced Very High Resolution Radiometer (AVHRR). The measured quantity is the sea surface radiance which is converted to an equivalent blackbody temperature. Correction factors are applied to account for atmospheric attenuation and radiation from atmospheric gases.

The infrared detectors have a temperature response from approximately  $-113^{\circ}$  to  $+47^{\circ}\text{C}$  (Brower et al., 1976). This is displayed in gray tones on photographic film, where the darker shades represent higher temperatures. Since the ocean temperature range is considerably smaller than the detector's range it is possible, through the technique of "image enhancement," to assign the available gray shades to this narrower range. This allows finer resolution of the gray shades that represent the oceanic temperature structure and will better distinguish thermal features.

Cloud cover is often difficult to distinguish in infrared images. In this case a comparison between the visual



the ratio of nitrate to phosphate is not very often 16:1 in ocean surface waters (Banse, 1974).

### III. METHODS

#### A. NUTRIENTS

Surface nutrient concentrations were determined approximately every 0.6 km except for the 13 June Cruise Leg 3 when engine problems caused speed adjustments. Seawater samples were pumped from a keel intake at 2.5m to the ship-board laboratory. Nitrate and phosphate were analyzed every two minutes according to the Technicon Autoanalyzer Industrial Method 175-72-WM, 177-72-WM (Anonymous, 1973) and 100-70-WM (Anonymous, 1978). Here nitrate may include traces of in situ nitrite, since the nitrate is reduced to nitrite before measurements. However, according to Paulson (1972) there is little or no interference from surface nitrate in this area.

#### B. SATELLITE IMAGERY

The pre-cruise procedure was initiated approximately 2-3 weeks prior to the estimated departure time. Mr. Breaker of the National Environmental Satellite Service (NESS) at Redwood City, California, monitored and enhanced images from the TIROS-N satellite series. Several days prior to the cruise, the day of the cruise and sometimes during the cruise updated satellite information was passed via telephone. Information, re IR inferred thermal features, such as orientation, approximate center, size, and spatial relationship

relative to prominent landmarks proved invaluable in locating features and planning sampling strategy. Additionally, the direction of a feature's general movement and large-scale changes in shape may be inferred by viewing a time sequence of images.

### C. COMPUTATIONS

A linear regression analysis was performed and mean values were calculated for nitrate, phosphate and temperature using the NORLSQ library subroutine at the W. R. Church Computer Center of the Naval Postgraduate School.

Correlation coefficients for nitrate to phosphate, nitrate to temperature and phosphate to temperature were obtained utilizing the correlation coefficient equation:

$$r = \Sigma(X_i - \bar{X})(Y_i - \bar{Y}) / ([\Sigma(X_i - \bar{X})^2][\Sigma(Y_i - \bar{Y})^2])^{1/2}$$

Zero values of nitrate and phosphate occurred when taking reference baselines or calibrating standards, or when a nitrate "none detected" condition occurred due to concentrations below the sensitivity of the instrument. When grossly anomalous values occurred, the nutrient and temperature data for that time increment were not used in the calculations.

(Statistical computations for the wind stress, adenosine triphosphate (ATP), and chlorophyll biomass will be discussed in a related thesis.)

#### D. CHLOROPHYLL

Fluorescence was recorded continuously using a Turner Model 111 fluorometer as described by Lorenzen (1966). Discrete samples, taken every 30 minutes, were used to calibrate and convert fluorescence to chlorophyll a concentration. The discrete chlorophyll samples were analyzed using the method of Strickland and Parsons (1968). These samples were filtered through a Whatman 4.5 cm G/FC glass fiber filter. For comparison with ATP, chlorophyll concentrations were converted to carbon units (mg/l) by using the average conversion factor (100) proposed by Holm-Hansen (1969).

#### E. ATP

Adenosine triphosphate was sampled every 10 minutes or approximately every 3 km over most of the cruise tracks. Each 50 ml sample was filtered through a 200- $\mu$ m nylon screen, then through a 0.45- $\mu$ m glass filter. The method of Holm-Hansen and Karl (1976) was used for ATP analysis. For comparison with chlorophyll, ATP concentrations were converted to carbon units (mg/l) by using the average conversion factor (250) proposed by Holm-Hansen (1969).

#### F. TEMPERATURE

Sea surface temperature was recorded continuously from a thermistor located at approximately 2.5 m depth coincident to the sea water intake. Sea surface temperature was also measured by bucket thermometer every 20 minutes simultaneously

with the release of a Sippican expendable bathythermograph probe (XBT). The bucket thermometer readings differed by a maximum of  $\pm 0.5^{\circ}\text{C}$  with respect to the continuous samples and by a maximum of  $\pm 0.6^{\circ}\text{C}$  with respect to the XBT temperatures. When these deviations exceeded  $\pm 0.3^{\circ}\text{C}$  they were adjusted to the bucket thermometer readings.

In Figures 8 and 21 the XBT data are plotted as isotherms in vertical sections along the cruise tracks for the 30 April and 7 August cruises. A contour interval of  $0.5^{\circ}\text{C}$  was used.

#### G. WIND

Wind effects were calculated from the Fleet Numerical Oceanography Center's (FNOC) Data Base using the so-called Field By Information Blending (FIB) routine for grid position  $36^{\circ}\text{N} \times 122^{\circ}\text{W}$ , which is in the vicinity of the Point Sur coastal region. Computer programs developed by Mr. Andrew Bakun of the National Marine Fisheries Service (Bakun, 1973) produced an output consisting of six hour values for the wind vector, wind stress magnitude, Ekman transport vector, upwelling index (the Ekman vector component normal to the coastline) and vertical velocity (Appendix B). A minimum of 12 days of data prior to and after each cruise was selected for analysis.

#### IV. RESULTS

##### A. 30 APRIL CRUISE

Satellite IR imagery indicated a cyclonic thermal feature had developed off Pt. Sur, California, on 19 April 1979 (Plate 1). It appeared to be forming as a seaward extension of coastally upwelled water. By 29 April it appeared to have persisted as a plume-like feature with a cyclonic swirl (Plate 2). Through the use of temperature and nutrient sampling, referenced to the satellite imagery, the central area was found on the first transect and re-located on each subsequent transect (Fig. 1).

Temperature, nitrate and phosphate are plotted against elapsed distance from  $36^{\circ}39.1'N$  x  $121^{\circ}58.6'W$  (Fig. 2). The feature is evident in the gradients of nitrate, phosphate and temperature.

Correlation coefficients of  $r = 0.98$  for nitrate to phosphate,  $r = -0.94$  for phosphate to temperature, and  $r = -0.96$  for nitrate to temperature were obtained (Table I). Nutrient ratio (N/P) and temperature when plotted against elapsed distance (Fig. 4) also exhibit a strong negative correlation with respect to each other; as is to be expected.

Based on the temperature minima and nutrient maxima the central area of the feature appeared at approximately 65 km (Leg 1), 220 km (Leg 2) and 300 km (Leg 3). The other cold water areas at approximately 165 km (Leg 2) and beyond 335 km





Plate 1. TIROS-N Satellite IR Image of the California Coast, 19 April 1979.




Plate 2. TIROS-N Satellite IR Image of the California Coast, 29 April 1979.




Fig. 1. Track of the 30 April Cruise (solid line) and of the Upwelling Feature (dashed line) based on Satellite IR Imagery, Sea Surface Temperature and Nutrient Data.




Fig. 2. Nitrate, Phosphate, and Sea Surface Temperature Versus Elapsed Distance along the Track of the 30 April Cruise. Note the tendency for inverse correlation between temperature and nutrients.




Fig. 3. ATP, Chlorophyll a, and Sea Surface Temperature Versus Elapsed Distance along the Track of the 30 April Cruise.




Fig. 4. Nutrient Ratio and Sea Surface Temperature Versus Elapsed Distance along the Track of the 30 April Cruise. Note the tendency for inverse correlation between temperature and nutrient ratio.




Fig. 5. Nitrate Versus Phosphate for the 30 April Cruise.




Fig. 6. Nitrate Versus Temperature for the 30 April Cruise.






Fig. 7. Phosphate Versus Temperature for the 30 April Cruise.




Fig. 8. Vertical Temperature Sections Along the Track of the 30 April Cruise.

(Leg 3) occurred when the ship's track extended into the near-shore upwelling region. Sharp chemical and thermal gradients or "fronts" are evident on all legs.

The linear regression analysis of nitrate versus phosphate, nitrate versus temperature and phosphate versus temperature (Figs. 5, 6, 7) yielded slopes of 15.10, -6.48 and -0.43 respectively and x-axis intercepts of 0.48  $\mu\text{M}$ , 12.99°C and 14.11°C respectively (Table I).

Figure 8 gives some idea of the vertical structure of the upwelled feature. The thermal feature is evident from elapsed distance 64.0 through 81.6 km, 206.2 through 213 km and 312.4 through 323.4 km (where the 11.0°C isotherm surfaces). Coastal upwelling is evident from elapsed distance 163.7 through 171.4 km and 329.3 through 353.8 km.

#### B. 13 JUNE CRUISE

On 13 June 1979, Mr. Laurence Breaker (NESS, Redwood City) gave approximate coordinates and adjacent shore features of a coastal upwelling event off Point Sur, with "plume-like" characteristics (Plate 3). The ACANIA sailed south parallel to the shore until the northern and southern thermal and nutrient gradients were transected. North-south legs were to be repeated westward in a ladder-like fashion until the seaward termination of the plume was found. However, the cruise was terminated (after only ten hours) due to a clogged sampling port.

Sea surface temperature, nutrient, ATP and chlorophyll data were collected along one and one-half transects of the feature as shown in Fig. 9. The elapsed distances are from point 36°36.3'N x 121°58.7'W.

Based on the distributions of temperature, nitrate and phosphate (Fig. 10) the upwelling area was encountered at approximately 15 km along the track. Nitrate and phosphate values were high (up to 12.1  $\mu$ M and 1.23  $\mu$ M respectively) until the southern frontal boundary was reached. In the oceanic water, at approximately 50 km, nitrate and phosphate concentrations were low (0.55 to 3.8  $\mu$ M and 0.33 to 0.65  $\mu$ M, respectively). To the north, the frontal southern boundary was again encountered at approximately 60 km along the track. Throughout, there was a strong correlation ( $r = 0.93$ ) between nitrate and phosphate, a strong inverse correlation between nitrate and temperature ( $r = -0.92$ ) and a strong inverse correlation between temperature and phosphate ( $r = -0.93$ ).

The chlorophyll data are unreliable after an elapsed distance of 40 km, because of excessive air injection into the sea chest during rough seas. Additionally, the XBT recorder was inoperative during this cruise.

#### C. 7 AUGUST CRUISE

The satellite image of 30 July 1979 (Plate 4) shows a cold water plume extending approximately 150 km southwest from Point Sur. Mr. Laurence Breaker (NESS, Redwood City)





Plate 3. TIROS-N Satellite IR Image of the California Coast, 13 June 1979.



CRUISE VI 13 JUNE, 1979

LEG 1 2100-2318 GMT

LEG 2 2318-2350 GMT

LEG 3 2350-0132 GMT

Fig. 9. Track of the 13 June Cruise (solid line) and Outline of the Upwelling Feature (dashed line) Based on Satellite Imagery, Sea Surface Temperature and Nutrient Data.




Fig. 10. Nitrate, Phosphate, Nutrient Ratio, ATP, Chlorophyll a, and Sea Surface Temperature Versus Elapsed Distance along the Track of the 13 June Cruise.




Fig. 11. Nitrate Versus Phosphate for the 13 June Cruise.






Fig. 12. Nitrate Versus Temperature for the 13 June Cruise.




Fig. 13. Phosphate Versus Temperature for the 13 June Cruise.

reported the feature had continued to extend southwest. By 5 August 1979 the feature had extended approximately 240 km southwest of Point Sur (Plate 5). Sea surface temperatures were expected to be within approximately 14 to 15°C inside the plume and greater than 17°C outside the plume based on ship reports correlated to the satellite image by Mr. Breaker.

The cruise plan was similar to the 30 April and 13 June cruises except, in this case, the primary interest was the water near the seaward termination of the feature. The track strategy was planned to coincide with the axis of the feature; when applicable, turn south to transect its southern boundary, zigzag across its southern boundary, then run back up the axis upon return. Due to limited sampling resources, ATP sampling did not commence until the southern boundary was reached.

The cruise track is shown in Fig. 14. Legs 2 through 5 intersected the feature's southern boundary. Elapsed distances are measured from position 36°38.8'N x 121°57.5'W.

Figure 15 is the line graph of temperature, nitrate and phosphate along the cruise track. The nutrient lines are discontinuous from 160 to 200 km, 275-295 km and 325-355 km because the Autoanalyzer was secured for cleaning. When there is a phosphate line but no line for nitrate, nitrate concentrations were so low that they were below the sensitivity of the instrument. However, in all cases the frontal crossings are clearly evident in the temperature lines.

Frontal crossings occurred at approximately 260, 325, 360 and 385 km on Legs 2 through 5. The line graphs for Legs 1 and 6 are generally within a temperature range of 14 to 16°C, indicating that the ACANIA was within the plume.

The correlation coefficients were  $r = 0.42$  for nitrate versus phosphate,  $r = -0.11$  for nitrate versus temperature, and  $r = -0.39$  for phosphate versus temperature. Linear regression analysis yielded slopes of 26.44, -13.63 and -0.32 and x-axis intercepts of 0.61  $\mu\text{M}$  phosphate, 15.35°C and 17.56°C for Figs. 18, 19, and 20, respectively (Table I).

The thermal feature is evident in the vertical cross section of temperature (Fig. 21) and gives some idea of the vertical structure of the feature. The transition between plume water and oceanic water is located between elapsed distances 249.5 through 263.1 km, 274.8 through 285.9 km, and 305.2 through 336.0 km (where the 15.5°C isotherm rises toward the surface). The 15.5°C isotherm is 30 to 40 meters deep outside the plume and inside it rises toward the surface. The thermal patchiness, evident in the satellite IR imagery, is seen in both the temperature cross section and the surface temperature line graphs.




Plate 4. TIROS-N Satellite IR Image of the California Coast, 30 July 1979.




Plate 5. TIROS-N Satellite IR Image of the California Coast, 5 August 1979.




Fig. 14. Track of the 7 August Cruise (solid line) and Outline of the Upwelling Feature (dashed line) based on Satellite Imagery, Sea Surface Temperature and Nutrient Data.




Fig. 15. Nitrate, Phosphate, and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise.






Fig. 16. ATP, Chlorophyll *a*, and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise.




Fig. 17. Nutrient Ratio and Sea Surface Temperature Versus Elapsed Distance Along the Track of the 7 August Cruise Track.




Fig. 18. Nitrate Versus Phosphate for the 7 August Cruise.




Fig. 19. Nitrate Versus Temperature for the 7 August Cruise.




Fig. 20. Phosphate Versus Temperature for the 7 August Cruise.




Fig. 21. Vertical Temperature Sections Along the Track of the 7 August Cruise

TABLE I  
SUMMARY OF REGRESSION ANALYSES

CRUISE	N ( $\mu\text{M}$ )	$\bar{P}$ ( $\mu\text{M}$ )	$\bar{T}$ ( $^{\circ}\text{C}$ )	SLOPE			"X" INTERCEPT			CORRELATION COEFFICIENTS		
				$\Delta\text{N}:\Delta\text{T}$	$\Delta\text{N}:\Delta\text{T}$	$\Delta\text{N}:\Delta\text{P}$	N:T( $^{\circ}\text{C}$ )	P:T( $^{\circ}\text{C}$ )	N:P( $\mu\text{M}$ )	N:T	P:T	N:P
April 30 to May 1	8.59	1.03	11.7	-6.48	-0.43	15.1	12.99	14.11	0.48	-0.96	-0.94	0.98
							13.79	14.43	0.26	-0.92	-0.93	0.93
June 13	8.24	0.90	12.4	-4.99	-0.39	12.88	15.35	17.56	0.61	-0.11	-0.39	0.42
August 7 to 9	5.28	0.767	15.06	-13.63	-0.32	26.44						

TABLE II  
DATA RANGES

TEMPERATURE °C	NITRATE μM	PHOSPHATE μM	RATIO N/P	ATP ng/l	CHLOROPHYLL mg/m <sup>3</sup>	ATP mgC/l	CHLOROPHYLL mgC/l
<u>30 April to 1 May Cruise</u>							
9.4 to 13.6	0.29 to 23.11	0.45 to 2.06	0.6:1 to 11.7:1	338 to 4126	0.13 to 6.49	0.0845 to 1.0315	0.013 to 0.649
<u>13 June Cruise</u>							
11.1 to 13.8	0.55 to 12.10	0.33 to 1.29	1.7:1 to 12.3:1	825 to 2485	1.15 to 17.28	0.206 to 0.621	0.115 to 1.728*
<u>7 to 9 August Cruise</u>							
12.2 to 17.2	none detected to 20.91	0.03 to 2.07	0.0 to 15.3:1	712 to 3418	0.17 to 8.19	0.178 to 0.855	.017 to 0.819

\*Data are unreliable due to air injection into the sampling water.



## V. DISCUSSION

A better understanding of the "life cycle" of an upwelling event may be provided through the integration of satellite IR imagery and in situ sampling. This approach leads to a descriptive "phase" model of the formation and dissipation of an upwelling chemical system and some insight on the interaction of dynamic and biological processes in the system. This thesis analyzes results obtained while using this integrated technique to investigate three upwelling events.

An upwelling event, like many physical phenomena, may be described in terms of phases: initiation, growth, equilibrium, decay and dissipation. Each phase is a function of the magnitude and rate of change of the dynamic processes (mixing, advection, diffusion) and the boundary conditions (atmospheric forcing; topography and bathymetry). The chemical characteristics of any phase of an upwelling event are a function of the initial conditions (source water); the magnitude, direction, and duration of the driving function (wind stress); and the differential utilization and regeneration by biological processes.

The satellite imagery prior to all cruises show "plumes" of colder water extending from the coast near Point Sur. These features range in scale from tens of kilometers to several hundred kilometers in length. Analysis of in situ

surface nutrient concentrations, summarized in Tables I and II, the surface winds in the area prior to and during the cruises (Appendix B), sea surface temperatures, satellite IR imagery and bathymetry strongly argue that these satellite detected thermal plumes were caused by upwelling events. That nutrient concentrations may closely follow the pattern of the satellite thermal imagery of the sea surface was shown by Traganza et al. (1980). These upwellings of cold nutrient-rich waters produce chemical and thermal fronts which appear to have associated with them increased biological activity.

The feature examined during the 30 April Cruise is postulated to be composed of recently upwelled water. Figure 2 shows very sharp chemical (up to  $\frac{10.41 \mu\text{M NO}_3}{4.7 \text{ km}}$ ) and thermal (up to  $0.9^\circ\text{C}/2.3 \text{ km}$ ) gradients. The range of nutrient ratio values are 0.6N:1P (lowest value, outside the feature) and 11.7N:1P (maximum level, inside the feature). The graph of nutrient ratio (Fig. 4) against elapsed distance along the track shows that the concentrations of high values are either within the area of the feature or the coastal upwelling. The low values of temperature (below  $12^\circ\text{C}$ ) and the high concentrations of nitrate and phosphate (up to  $12.1 \mu\text{M}$  and  $1.29 \mu\text{M}$ , respectively) are all either within the feature or the coastal upwelling (Fig. 10). There are strong correlations between nitrate and phosphate ( $r = 0.98$ ), nitrate and temperature ( $r = 0.96$ ), and phosphate and temperature ( $r = -0.94$ ). Analysis of the wind data prior to the

cruise shows that favorable upwelling conditions occurred during the periods 9 to 20 April and 28 April to 4 May (Appendix B). All these factors, plus the lower temperatures indicated by the satellite IR imagery are taken to describe a feature of recently upwelled water in the early phases of its life cycle.

The 13 June Cruise examined a feature that is close inshore (Plate 3). Wind data (Appendix B) shows favorable upwelling conditions commenced only three days prior to cruise. The waters outside the feature are about 14°C but inside the feature temperatures fall below 12°C. All high concentrations of nutrients are within the feature, as are all high nutrient ratios. There are strong correlations between nitrate and phosphate ( $r = 0.93$ ), nitrate and temperature ( $r = -0.92$ ), and phosphate and temperature ( $r = -0.93$ ) (Figs. 11, 12 and 13). This feature is recently upwelled water and is postulated to be in the early phases of the upwelling event "life cycle."

The waters examined in these two cruises still have very strong source water (thermal and nutrient) signatures. The dynamic and biological processes have not had sufficient time to change the upwelled water's initial characteristics. Along the frontal boundaries, the biological processes become evident (Figs. 3 and 10). The dynamic processes appear to have provided the necessary environment for biological activity.

The data from the 7 August Cruise were gathered predominantly from within the feature (Fig. 14). In comparison to the previous cruises, this resulted in a considerably smaller proportion of data points being collected outside the feature. A statistically insufficient number of samples were obtained in the low nutrient concentration, high temperature region. In comparison to the previous cruises the analysis is biased as follows: mean nutrient concentrations are high; mean temperature is low; the slopes of nitrate to phosphate, nitrate to temperature, and phosphate to temperature are too high, and the respective x-axis intercepts are too low.

The feature observed on the 7 August Cruise appears to contain upwelled water that is in various phases of the upwelling cycle. At 50 km elapsed distance water temperature was below 13°C with rapidly increasing nutrient ratios. Sampling was interrupted to clean the Autoanalyzer but concentrations were approaching levels consistent with the previous cruises (nutrient ratios above 10N:1P). This area of the feature contains water that has source water characteristics and may be assumed to be in the early phases. At 75 km elapsed distance, while still within the feature, temperature was approximately 14°C and nutrients had fallen, but not to the "outside-the-features-levels" of the previous cruises. This area might be considered to be close to equilibrium. The water was still relatively cold and nutrient-rich but no longer similar to the source water; the

dynamic and biological processes apparently had started to modify it. The temperature then rose to the 15.0° to 15.5°C range and while nutrients still generally decreased with increasing temperature, they showed considerable variability. This area can be characterized as in the decay phase. The dynamic and biological processes continued to change the temperature and nutrient characteristics.

Once outside the feature, temperatures rose sharply to levels of 17.5°C and nutrients correspondingly fell. As the track moved west (Fig. 15) the nutrient levels and nutrient ratios (Fig. 17) never recovered to the levels previously encountered within the feature. In some areas, at the western most extent of the cruise, the nitrate levels are too low to be detected. In most coastal upwelling systems, nitrogen is characteristically the limiting nutrient, Dugdale et al. (1967) and Thomas, (1969). "In accordance with Liebig's law of the minimum, that constituent of the sea water present in smallest quantity relative to the requirement for growth of organisms will become the limiting factor." (Redfield, 1958). In this area, there is insufficient nitrate to sustain photosynthesis and, therefore, the biological modification of the feature is complete and low biomass concentrations are evident (Fig. 16). All that remains is for the dynamic processes to dissipate the temperature anomaly.

The wind data prior to the cruise shows a cyclical pattern of upwelling events occurring since early April.

Upwelling periods were of the order of one to three weeks and relaxation periods were of the order of two to six days (Appendix B). The average Ekman transport varied from west southwest to southwest. With such an intermittent, long range pattern of offshore transport; it is therefore possible that the feature investigated on the 7 August Cruise was in fact upwelled water of varying ages. Barton, et al. (1977) indicates that during an upwelling event coldest water will migrate to the continental shelf edge and remain there until the system relaxes. Upon return of favorable winds the upwelling will again reappear along the inner shelf. In this case, high nutrient, cold water was found at 50 km elapsed distance which is just inside the shelf break. The center of an upwelling may stop at the shelf edge, with continued favorable winds, upwelled water will continue to be transported offshore in a 20 m thick surface layer (Huyer, 1974). Patchiness or banding can result from this start/stop action between close upwelling events (Barton, et al. 1977). The satellite images (Plates 4 and 5) show some patchiness. The second image (Plate 5) shows the feature has moved 90 km further southwest and the wind data show the Ekman transport to be west southwesterly during the period between the images. Therefore the feature investigated has some of the characteristics of a wind induced upwelling plume with the oldest water at its furthest extent; still maintaining some of its temperature characteristics but due to biological action and mixing losing much of its high nutrient signature at its westernmost (oldest) edges.

In viewing the correlation values of all cruises it must be remembered "that where elements are substantially depleted by the growth of plants small unused residues of one or another element may greatly alter the ratios." (Redfield, 1958). The nutrient-to-temperature correlations will reflect the effects of heat transfer, mixing, advection, diffusion and biological utilization and regeneration. The longer the water has been on the surface, the more time the processes have to affect the correlations. This is evident on the 7 August Cruise where all correlations are poor.

In sea water the ratio of change of nitrate and phosphate,  $\Delta N/\Delta P$ , is representative of the uptake of these nutrients by phytoplankton at a rate of 16N:1P. The slope of the best fit line found from linear regression analysis (Table 1) represents this ratio of change. The values obtained on the 30 April Cruise (15.1:1) and 13 June Cruise (12.88:1) are close to the theoretical value of 16N:1P.

On all cruises the nutrient ratio did not equal the theoretical value of 16:1 (Redfield, 1958) most likely because the source water was not at that ratio. The ratios ranged from a low where no nitrate was detected to a high of 15.3:1. The low values perhaps were found where further biological uptake had become inhibited due to low nitrate concentration and the higher values corresponded to the upwelled water. These values are not inconsistent with the annual range for the nutrient ratio of 3:1 to 13:1 obtained in a study by Butler et al. (1979).

Finally, in areas of upwelling there exist some geographically fixed preferred positions, e.g. south of capes, where plumes of freshly upwelled water protrude offshore (Shaffere, 1976; Reid et al., 1958). The research of this thesis and that of Traganza et al. (1980), which covered one annual cycle, indicates the coastal waters off Point Sur, California, are such an area. Whether this is because of the proximity of Point Sur and/or Monterey Canyon, local bathymetry, local water circulation patterns, local wind effects, or coastally trapped topographic Rossby waves; or a combination of these factors cannot be determined from the data. However, offshore plumes consistently, albeit aperiodically, originate from this area.



## VI. CONCLUSIONS

1. High resolution, enhanced satellite IR imagery can be of great value in locating and assessing feature motion when investigating upwelling events that are manifested as sea surface temperature anomalies.
2. Upwelled water may be evidenced at considerable (ca. 100 to 300 km) distances from the coast.
3. Qualitative inferences on the distribution of nutrient concentrations can be made using satellite infrared imagery if coupled with in situ sampling to establish nutrient versus temperature correlations.
4. Altogether, with sufficient historical upwelling data from in situ monitoring, a nutrient predictive model for upwelled water in the initiation, growth and equilibrium phases is conceivable in the future.
5. A predictive model of the entire upwelling cycle, in all its phases, is not yet possible. The effects of air-sea interaction, the dynamical processes, and the biological processes make any nutrient model extremely complex. The dynamic processes vary from region to region and season to season. Regeneration and utilization rates vary from organism to organism and season to season. This further compounds the task.
6. The area off Point Sur, California, consistently has offshore upwelling plumes and merits further detailed

study. Such study is continuing under the direction of Dr. Eugene Traganza supported by the Office of Naval Research, Code 482.

7. From a Naval aspect the thermal gradients in upwelled water can have a considerable impact on sound propagation. The increased biological concentrations associated with the thermochemical fronts can, depending on the species, increase the reverberation levels and background noise.

# APPENDIX A

R/V ACANIA										30 APR / 1 MAY 79	CHEMICAL MESOSCALE CRUISE VI)	
TIME GMT 1604	LATITUDE N 36 39.1	LONGITUDE WEST 121 58.6	DISTANCE KM 0.0	NP3 NM 0.48	PN4 GM 0.53	MUTR-RATIO NO3/PN4 0.9	ATP NG/L	CHL A MG/23 5.57	TEMP C 12.6			
			4.7	0.83	0.56	1.5		5.66	12.6			
			5.8	0.67	0.76	4.8	2668.0	4.34	12.3			
			6.4	1.36	0.61	2.2		4.35	12.3			
			7.0	0.83	0.55	1.5		4.18	12.3			
			8.2					4.29	12.4			
			9.3					4.42	12.4			
			9.9					4.66	12.4			
			10.5	1.54	0.56	2.8		4.71	12.4			
			11.1	2.60	0.65	4.0		5.04	12.4			
			11.7	5.62	0.86	6.5		4.73	12.0			
			12.3	7.21	0.96	7.7		4.64	11.7			
			13.4	7.15	1.01	7.7		3.53	11.5			
			14.0	10.41	1.18	8.8	2046.0	2.59	11.2			
			15.2	9.16	1.19	7.8		1.57	11.2			
			16.3	10.23	1.20	8.5		1.39	11.2			
			17.5	10.58	1.21	8.4		1.16	11.3			
			18.1	10.58	1.22	8.7		1.01	11.3			
			19.3	10.23	1.23	8.6		1.02	11.3			
			19.9	10.05	1.20	8.5		1.39	11.3			
			20.4		1.19	8.4	2299.0	1.41	11.4			
			21.0				3303.0	1.17	11.5			
			22.2	6.86	0.92	7.5		3.43	11.5			
			23.3	7.57	0.95	8.0		3.64	11.5			
			24.4	7.04	0.93	8.0		3.43	11.5			
			25.5	6.80	0.92	7.6	2534.0	3.13	11.9			
			26.6	6.15	0.89	6.9		3.16	11.9			
			27.4	5.62	0.86	6.7		3.14	12.0			
			28.0	5.97	0.84	6.7	2060.0	3.74	12.0			
			29.2	5.97	0.86	6.9		3.11	12.1			
			30.3	6.15	0.89	6.9	2170.0	3.50	12.1			
			31.5	6.50	0.92	7.0		3.08	12.1			
			32.7		0.93			3.07	12.1			

CHEMICAL MISCELLANEA (CRUIS = VI)

TRWD  
DEC 1  
12-0

● ●  
● ●

[illegible]

[illegible]

R/V ACANIA

30 APR / 1 MAY 79

CHEMICAL MESOSCALE CRUISE VII

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR. RATIO NO3/PO4	ATD NG/L	CHL A MG/WT	TEMP DEG C
2400	35 49.4	121 46.2	96.3	1.87	0.56	3.3		1.43	12.8
			97.4	1.87	0.56	3.3		1.43	12.8
			98.0					1.43	12.8
			98.6					1.43	12.8
			99.2					1.43	12.8
			99.8					1.43	12.8
			100.4					1.43	12.8
			101.0					1.43	12.8
			101.5					1.43	12.8
			102.1					1.43	12.8
			102.7					1.43	12.8
			103.3					1.43	12.8
			103.9	3.31	0.45	7.4		1.43	12.8
			104.4	0.0	0.45	0.0		1.43	12.8
0100	35 42.4	121 38.5	105.0	0.0	0.47	0.0		1.43	12.8
			105.6	0.0	0.47	0.0		1.43	12.8
			106.2	0.0	0.45	0.0		1.43	12.8
			106.8	0.0	0.44	0.0		1.43	12.8
			107.4	0.0	0.42	0.0		1.43	12.8
			107.9	0.0	0.44	0.0		1.43	12.8
			108.5	0.0	0.47	0.0		1.43	12.8
			109.1	0.0	0.46	0.0		1.43	12.8
			109.7	0.0	0.44	0.0		1.43	12.8
			110.3	0.0	0.45	0.0		1.43	12.8
			110.9	0.0	0.45	0.0		1.43	12.8
			111.4	0.0	0.44	0.0		1.43	12.8
			112.0	0.0	0.48	0.0		1.43	12.8
			112.6	0.0	0.54	1.2		1.43	12.8
			113.2	0.0	0.56	1.8		1.43	12.8
			113.8	1.73	0.62	2.8		1.43	12.8
			114.4	1.73	0.61	2.8		1.43	12.8
			115.0	1.73	0.61	2.8		1.43	12.8
			115.6	1.73	0.63	2.7		1.43	12.8
			116.2	1.73	0.63	2.7		1.43	12.8
			116.8	1.73	0.63	2.7		1.43	12.8
			117.4	1.73	0.63	2.7		1.43	12.8
			118.0	1.73	0.63	2.7		1.43	12.8
			118.6	1.73	0.63	2.7		1.43	12.8
			119.2	1.73	0.63	2.7		1.43	12.8
			119.8	1.73	0.63	2.7		1.43	12.8
			120.4	1.73	0.63	2.7		1.43	12.8
			121.0	1.73	0.63	2.7		1.43	12.8
			121.6	1.73	0.63	2.7		1.43	12.8
			122.2	1.73	0.63	2.7		1.43	12.8
			122.8	1.73	0.63	2.7		1.43	12.8
			123.4	1.73	0.63	2.7		1.43	12.8
			124.0	1.73	0.63	2.7		1.43	12.8
			124.6	1.73	0.63	2.7		1.43	12.8
			125.2	1.73	0.63	2.7		1.43	12.8
			125.8	1.73	0.63	2.7		1.43	12.8
			126.4	1.73	0.63	2.7		1.43	12.8
			127.0	1.73	0.63	2.7		1.43	12.8
			127.6	1.73	0.63	2.7		1.43	12.8
			128.2	1.73	0.63	2.7		1.43	12.8
			128.8	1.73	0.63	2.7		1.43	12.8
			129.4	1.73	0.63	2.7		1.43	12.8
			130.0	1.73	0.63	2.7		1.43	12.8
			130.6	1.73	0.63	2.7		1.43	12.8
			131.2	1.73	0.63	2.7		1.43	12.8

R/V ACAAHA

30 APR / 1 MAY 79

CHEMICAL MESOSCALE (CRUISE VI)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	HM3	PC4	NUMR.RATIO H33/P04	ATP NG/L	CHL A MG/1003	TAP C
0200	35 36.8	121 31.7	125.4	1.01	0.58	1.7	1023.0	1.15	13.0
			126.0	1.19	0.60	2.0	1773.0	1.24	13.0
			127.2	1.19	0.62	1.9	1121.0	1.41	13.0
			128.4	1.01	0.61	1.7	1264.0	1.44	13.0
			129.5	1.01	0.61	1.7	1051.0	1.34	13.0
			130.1	0.83	0.62	1.6	1297.0	1.49	13.0
			131.3	0.47	0.53	0.9	1264.0	1.77	12.9
			131.9	0.65	0.53	1.2	1121.0	1.58	12.9
			132.4	0.47	0.50	0.9	1264.0	1.56	12.9
			133.0	0.29	0.49	0.6	1023.0	1.46	13.0
			133.6	0.0	0.49	0.0	1264.0	1.36	13.0
			134.2	0.0	0.49	0.0	1051.0	1.23	13.0
			134.8	0.0	0.49	0.0	1264.0	1.11	13.0
			135.4	0.0	0.51	0.0	1264.0	0.87	13.0
			136.5	0.0	0.51	0.0	1051.0	0.75	13.0
			137.1	0.0	0.51	0.0	1264.0	0.85	13.0
			137.7	0.0	0.51	0.0	1051.0	0.83	13.0
			138.3	0.0	0.51	0.0	1264.0	0.92	13.0
			139.0	0.0	0.51	0.0	1051.0	0.80	13.0
			140.0	0.0	0.51	0.0	1264.0	0.87	13.0
			141.2	0.0	0.51	0.0	1051.0	0.87	13.0
			142.4	0.0	0.51	0.0	1264.0	0.85	13.0
			143.5	0.0	0.51	0.0	1051.0	0.84	13.0
			144.1	0.0	0.51	0.0	1264.0	0.85	13.0
			144.7	0.0	0.51	0.0	1051.0	0.85	13.0
			145.3	0.0	0.51	0.0	1264.0	0.86	13.0
			146.0	0.0	0.51	0.0	1051.0	0.86	13.0
			147.0	0.0	0.51	0.0	1264.0	0.87	13.0
			148.2	0.0	0.51	0.0	1051.0	0.87	13.0
			149.4	0.0	0.51	0.0	1264.0	0.87	13.0
			149.9	0.0	0.51	0.0	1051.0	0.87	13.0
			150.5	0.0	0.51	0.0	1264.0	0.87	13.0
			151.1	0.0	0.51	0.0	1051.0	0.87	13.0
			152.3	0.0	0.51	0.0	1264.0	0.87	13.0
			153.4	0.0	0.51	0.0	1051.0	0.87	13.0
			154.0	0.0	0.51	0.0	1264.0	0.87	13.0





R/V ACANIA

30 APR / 1 MAY 79

CHEMICAL MESUSCALE (CRUISE VII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NU3 TIM	PM4 TIM	NUTR. RATIO NU3/PM4	ATP NG/L	CHL A MG/43	TEMP DEG. C
0500	35 57.7	121 41.8	183.8	5.88	0.85	6.9		3.35	12.1
			184.4	5.88	0.85	6.9		3.35	12.0
			185.0	5.83	0.87	6.8		3.31	11.9
			185.5	5.34	0.82	6.5		3.31	11.8
			186.1	4.62	0.79	5.8	919.0	3.31	11.6
			186.6	4.26	0.75	5.6		3.31	11.5
			187.3	4.08	0.74	5.5		3.31	11.4
			187.9	3.72	0.71	5.2	880.0	3.31	11.3
			188.5	3.54	0.71	5.0		3.31	11.2
			189.0	3.72	0.73	5.2		3.31	11.1
			189.6	4.08	0.75	5.4		3.31	11.0
			190.2	4.80	0.78	6.2		3.31	10.9
			191.4	4.26	0.73	5.7	1036.0	3.31	10.8
			192.0	4.26	0.73	5.7		3.31	10.7
0600	36 1.7	121 51.8	192.5	3.90	0.72	5.3		3.31	10.6
			193.1	3.72	0.72	5.2		3.31	10.5
			193.7	3.54	0.73	5.4	1070.0	3.31	10.4
			194.3	4.08	0.75	5.7		3.31	10.3
			194.9	4.62	0.81	6.5		3.31	10.2
			195.5	4.80	0.81	6.5		3.31	10.1
			196.0	4.44	0.78	5.9		3.31	10.0
			196.6	4.26	0.76	5.7	1194.0	3.31	9.9
			197.2	4.44	0.76	5.7		3.31	9.8
			197.8	4.26	0.76	5.6		3.31	9.7
			198.4	4.44	0.76	5.7		3.31	9.6
			199.0	4.62	0.81	6.5	4126.0	3.31	9.5
			199.5	4.44	0.78	5.7		3.31	9.4
			200.1	4.26	0.72	5.2		3.31	9.3
			200.7	2.81	0.72	3.9		3.31	9.2
0600	36 1.7	121 51.8	201.3	2.45	0.69	3.6		3.31	9.1
			201.9	2.45	0.69	3.6		3.31	9.0
			202.5	3.17	0.77	4.3	3583.0	3.31	8.9
			203.0	3.17	0.77	4.3		3.31	8.8
			203.6	3.17	0.77	4.3		3.31	8.7
			204.2	3.17	0.77	4.3		3.31	8.6
			204.8	3.17	0.77	4.3		3.31	8.5
			205.4	3.17	0.77	4.3	3247.0	3.31	8.4
			206.0	3.17	0.77	4.3		3.31	8.3
			206.6	3.17	0.77	4.3		3.31	8.2
			207.2	3.17	0.77	4.3		3.31	8.1
			207.8	3.17	0.77	4.3		3.31	8.0
			208.4	3.17	0.77	4.3	2952.0	3.31	7.9
			209.0	3.17	0.77	4.3		3.31	7.8
			209.6	3.17	0.77	4.3	2066.0	3.31	7.7

P/V ACANIA

30 APR / 1 MAY 79

CHEMICAL WTS/SCALE/CRUISE VI)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE NM	UHQ	UHQ	NUTR. RATIO NO3/PO4	ATP NG/L	CHL A MG/M3	TEMP DEG C
0700	36	5.2	122	1.6					
			213.0	13.92	1.30	10.2		1.96	11.3
			214.1	14.36	1.39	10.3		2.23	11.1
			214.7	15.26	1.44	10.6	2904.0	2.17	10.5
			215.3	15.44	1.44	10.6		2.61	10.5
			215.9	16.70	1.54	10.8		1.97	10.6
			216.5	17.61	1.60	11.0		1.54	10.4
			217.0	18.51	1.65	11.2		1.30	10.2
			217.6	18.87	1.66	11.2	811.0	1.44	10.2
			218.1	19.41	1.71	11.2		1.75	10.0
			218.8	19.77	1.71	11.2		1.74	9.9
			219.4	19.77	1.76	11.2		1.74	9.9
			220.0	19.41	1.76	11.2		1.74	9.9
			220.5	19.23	1.76	11.0	974.0	1.74	9.7
			221.1	19.07	1.72	11.1		1.96	9.7
			221.7	18.87	1.70	11.1		1.05	9.8
			222.3	18.09	1.71	11.1		1.97	10.0
			222.9	17.51	1.69	11.0	545.0	1.17	10.0
			223.5	17.15	1.68	10.8		1.15	10.0
			224.0	17.79	1.66	10.7		1.15	10.0
			224.6	17.79	1.66	10.8		1.15	10.1
			225.2	17.61	1.63	10.5	590.0	1.15	10.1
			225.8	17.61	1.63	10.7		1.15	10.2
			226.4	17.61	1.63	10.8		1.15	10.3
			227.0	17.61	1.63	10.5		1.15	10.5
			227.5	17.61	1.63	10.5		1.15	10.5
			228.1	17.61	1.63	10.5		1.15	10.5
			228.7	17.61	1.63	10.5		1.15	10.5
			229.3	17.61	1.63	10.5		1.15	10.5
			229.9	17.61	1.63	10.5		1.15	10.5
			230.5	17.61	1.63	10.5		1.15	10.5
			231.1	17.61	1.63	10.5		1.15	10.5
			231.6	17.61	1.63	10.5		1.15	10.5
			232.2	17.61	1.63	10.5		1.15	10.5
			232.8	17.61	1.63	10.5		1.15	10.5
			233.4	17.61	1.63	10.5		1.15	10.5
			234.0	17.61	1.63	10.5		1.15	10.5
			234.6	17.61	1.63	10.5		1.15	10.5
			235.2	17.61	1.63	10.5		1.15	10.5
			235.8	17.61	1.63	10.5		1.15	10.5
			236.4	17.61	1.63	10.5		1.15	10.5
			237.0	17.61	1.63	10.5		1.15	10.5
			237.6	17.61	1.63	10.5		1.15	10.5
			238.2	17.61	1.63	10.5		1.15	10.5
			238.8	17.61	1.63	10.5		1.15	10.5
			239.4	17.61	1.63	10.5		1.15	10.5
			240.0	17.61	1.63	10.5		1.15	10.5
			240.6	17.61	1.63	10.5		1.15	10.5
			241.2	17.61	1.63	10.5		1.15	10.5
			241.8	17.61	1.63	10.5		1.15	10.5
			242.4	17.61	1.63	10.5		1.15	10.5
			243.0	17.61	1.63	10.5		1.15	10.5
			243.6	17.61	1.63	10.5		1.15	10.5
			244.2	17.61	1.63	10.5		1.15	10.5
			244.8	17.61	1.63	10.5		1.15	10.5
			245.4	17.61	1.63	10.5		1.15	10.5
			246.0	17.61	1.63	10.5		1.15	10.5
			246.6	17.61	1.63	10.5		1.15	10.5
			247.2	17.61	1.63	10.5		1.15	10.5
			247.8	17.61	1.63	10.5		1.15	10.5
			248.4	17.61	1.63	10.5		1.15	10.5
			249.0	17.61	1.63	10.5		1.15	10.5
			249.6	17.61	1.63	10.5		1.15	10.5
			250.2	17.61	1.63	10.5		1.15	10.5
			250.8	17.61	1.63	10.5		1.15	10.5
			251.4	17.61	1.63	10.5		1.15	10.5
			252.0	17.61	1.63	10.5		1.15	10.5
			252.6	17.61	1.63	10.5		1.15	10.5
			253.2	17.61	1.63	10.5		1.15	10.5
			253.8	17.61	1.63	10.5		1.15	10.5
			254.4	17.61	1.63	10.5		1.15	10.5
			255.0	17.61	1.63	10.5		1.15	10.5
			255.6	17.61	1.63	10.5		1.15	10.5
			256.2	17.61	1.63	10.5		1.15	10.5
			256.8	17.61	1.63	10.5		1.15	10.5
			257.4	17.61	1.63	10.5		1.15	10.5
			258.0	17.61	1.63	10.5		1.15	10.5
			258.6	17.61	1.63	10.5		1.15	10.5
			259.2	17.61	1.63	10.5		1.15	10.5
			259.8	17.61	1.63	10.5		1.15	10.5
			260.4	17.61	1.63	10.5		1.15	10.5
			261.0	17.61	1.63	10.5		1.15	10.5
			261.6	17.61	1.63	10.5		1.15	10.5
			262.2	17.61	1.63	10.5		1.15	10.5
			262.8	17.61	1.63	10.5		1.15	10.5
			263.4	17.61	1.63	10.5		1.15	10.5
			264.0	17.61	1.63	10.5		1.15	10.5
			264.6	17.61	1.63	10.5		1.15	10.5
			265.2	17.61	1.63	10.5		1.15	10.5
			265.8	17.61	1.63	10.5		1.15	10.5
			266.4	17.61	1.63	10.5		1.15	10.5
			267.0	17.61	1.63	10.5		1.15	10.5
			267.6	17.61	1.63	10.5		1.15	10.5
			268.2	17.61	1.63	10.5		1.15	10.5
			268.8	17.61	1.63	10.5		1.15	10.5
			269.4	17.61	1.63	10.5		1.15	10.5
			270.0	17.61	1.63	10.5		1.15	10.5
			270.6	17.61	1.63	10.5		1.15	10.5
			271.2	17.61	1.63	10.5		1.15	10.5
			271.8	17.61	1.63	10.5		1.15	10.5
			272.4	17.61	1.63	10.5		1.15	10.5
			273.0	17.61	1.63	10.5		1.15	10.5
			273.6	17.61	1.63	10.5		1.15	10.5
			274.2	17.61	1.63	10.5		1.15	10.5
			274.8	17.61	1.63	10.5		1.15	10.5
			275.4	17.61	1.63	10.5		1.15	10.5
			276.0	17.61	1.63	10.5		1.15	10.5
			276.6	17.61	1.63	10.5		1.15	10.5
			277.2	17.61	1.63	10.5		1.15	10.5
			277.8	17.61	1.63	10.5		1.15	10.5
			278.4	17.61	1.63	10.5		1.15	10.5
			279.0	17.61	1.63	10.5		1.15	10.5
			279.6	17.61	1.63	10.5		1.15	10.5
			280.2	17.61	1.63	10.5		1.15	10.5
			280.8	17.61	1.63	10.5		1.15	10.5
			281.4	17.61	1.63	10.5		1.15	10.5
			282.0	17.61	1.63	10.5		1.15	10.5
			282.6	17.61	1.63	10.5		1.15	10.5
			283.2	17.61	1.63	10.5		1.15	10.5
			283.8	17.61	1.63	10.5		1.15	10.5
			284.4	17.61	1.63	10.5		1.15	10.5
			285.0	17.61	1.63	10.5		1.15	10.5
			285.6	17.61	1.63	10.5		1.15	10.5
			286.2	17.61	1.63	10.5		1.15	10.5
			286.8	17.61	1.63	10.5		1.15	10.5
			287.4	17.61	1.63	10.5		1.15	10.5
			288.0	17.61	1.63	10.5		1.15	10.5
			288.6	17.61	1.63	10.5		1.15	10.5
			289.2	17.61	1.63	10.5		1.15	10.5
			289.8	17.61	1.63	10.5		1.15	10.5
			290.4	17.61	1.63	10.5		1.15	10.5
			291.0	17.61	1.63	10.5		1.15	10.5
			291.6	17.61	1.63	10.5		1.15	10.5
			292.2	17.61	1.63	10.5		1.15	10.5
			292.8	17.61	1.63	10.5		1.15	10.5
			293.4	17.61	1.63	10.5		1.15	10.5
			294.0	17.61	1.63	10.5		1.15	10.5
			294.6	17.61	1.63	10.5		1.15	10.5
			295.2	17.61	1.63	10.5		1.15	10.5
			295.8	17.61	1.63	10.5		1.15	10.5
			296.4	17.61	1.63	10.5		1.15	10.5
			297.0	17.61	1.63	10.5		1.15	10.5
			297.6	17.61	1.63	10.5		1.15	10.5
			298.2	17.61	1.63	10.5		1.15	10.5
			298.8	17.61	1.63	10.5		1.15	10.5
			299.4	17.61	1.63	10.5		1.15	10.5
			300.0	17.61	1.63	10.5		1.15	10.5
			300.6	17.61	1.63	10.5		1.15	10.5
			301.2	17.61	1.63	10.5		1.15	10.5
			301.8	17.61	1.63	10.5		1.15	10.5
			302.4	17.61	1.63	10.5		1.15	10.5
			303.0	17.61	1.63	10.5		1.15	10.5
			303.6	17.61	1.63	10.5		1.15	10.5
			304.2	17.61	1.63	10.5		1.15	10.5
			304.8	17.61	1.63	10.5		1.15	10.5
			305.4	17.61	1.63	10.5		1.15	10.5
			306.0	17.61	1.63	10.5		1.15	10.5
			306.6	17.61	1.63	10.5		1.15	10.5
			307.2	17.61	1.63	10.5		1.15	10.5
			307.8	17.61					

R/V ACANIA 30 APR / 1 MAY 79 CHEMICAL MESUSCALE (CRUISE VI)									
TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KN	NU3 UM	PO4 UM	NU3/PO4 RATIO	ATP MG/L	CHL A MG/23	FLC C
0800	36 9.1	122 12.6	242.1	3.17	0.67	4.7	684.0	0.65	12.0
			243.1	3.23	0.50	3.8		1.18	12.0
			243.9	1.87	0.35	5.4		0.69	12.4
			244.5	1.80	0.35	5.2		0.49	12.4
			245.0	1.12	0.52	2.2		0.75	12.8
			245.6	1.11	0.50	2.2		0.64	13.0
			246.2	0.91	0.50	1.8		0.53	12.9
			246.8	0.90	0.50	1.8		0.32	12.9
			247.4	1.07	0.50	2.1		0.15	12.8
			248.0	1.63	0.57	2.9		0.53	12.9
			248.5	2.18	0.57	3.8	635.0	0.53	12.9
			249.1	2.37	0.60	3.9		0.49	12.9
			249.7	2.35	0.60	3.9		0.52	12.9
			250.3	2.54	0.62	4.1		0.57	12.9
			250.9	2.54	0.62	4.1		0.37	12.7
			251.5	1.15	0.55	3.7		0.41	12.8
			252.1	1.76	0.55	3.2		0.41	12.8
			252.7	1.55	0.53	2.9		0.44	12.9
			253.2	1.54	0.53	3.0		0.45	12.9
			253.8	1.53	0.53	3.0		0.23	13.1
			254.4	1.71	0.53	3.1	577.0	0.23	13.0
			255.0	1.70	0.53	3.2		0.23	13.0
			255.6	2.68	0.59	4.5		0.22	13.1
			256.1	2.59	0.59	4.3		0.22	13.3
			256.7	2.67	0.59	4.5		0.22	13.3
			257.3	2.66	0.59	4.5		0.22	13.3
			257.9	2.66	0.58	4.6		0.19	13.0
			258.5	2.45	0.58	4.2		0.21	13.2
			259.1	2.45	0.58	4.2		0.21	13.2
			260.8	2.44	0.58	4.1		0.21	13.4
0900	36 12.2	122 23.2	261.4	2.32	0.56	3.8	446.0	0.37	13.4
			262.0	2.01	0.56	3.6		0.37	13.4
			262.6	2.01	0.56	3.6		0.37	13.4
			263.1	2.01	0.56	3.6		0.37	13.4
			263.7	1.79	0.56	3.2		0.54	13.0
			264.3	1.57	0.54	2.9		0.24	13.0
			264.9	1.36	0.54	2.5		0.24	13.2
			265.5	1.35	0.52	2.5		0.20	13.2
			266.1	1.34	0.53	2.5		0.20	13.2
			266.6	0.91	0.53	1.8		0.13	13.3
			267.2	0.90	0.48	1.9	342.0	0.15	13.3
			267.8	0.68	0.47	1.4		0.14	13.5
			268.4	0.47	0.47	1.0		0.14	13.6
			269.0	0.45	0.45	1.0		0.20	13.4
			269.6	0.44	0.45	1.0		0.32	13.4
			270.1	0.42	0.45	0.9		0.32	13.4
			270.6	0.38	0.45	0.9		0.32	13.4
			271.1	0.38	0.45	0.9		0.32	13.4
			271.6	0.38	0.45	0.9		0.32	13.4
			272.1	0.38	0.45	0.9		0.32	13.4
1100	35 52.9	122 14.3	313.9	9.38	1.03	9.1	513.0	2.06	14.5

R/V ACANIA

10 APR / 1 MAY 79

CHEMICAL METEOROLOGICAL CRUISE VIJ

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE NM	MDJ UM	PMJ UM	NOTE NUTR:RATIO NO3:PO4	ATP MG/L	CHL MG/3	TEMP C
1200	35	59.6	122	6.0					
			119.5	10.13	1.10	9.2		2.06	11.5
			115.1	10.50	1.13	9.3		2.06	11.5
			115.6	10.31	1.10	9.4		2.06	11.5
			116.2	10.13	1.10	9.3		2.06	11.5
			116.4	9.94	1.11	9.3		2.06	11.5
			117.4	9.42	1.10	7.4		1.87	11.4
			118.0	8.12	0.84	6.1		1.40	11.4
			119.1	6.09	0.99	8.2		1.37	11.4
			119.7	5.91	1.16	9.0		1.77	11.4
			120.3	5.91	1.16	9.2		1.77	11.4
			120.5	5.91	1.16	9.2		1.77	11.4
			121.5	5.91	1.16	9.2		1.77	11.4
			122.1	5.91	1.16	9.2		1.77	11.4
			122.6	5.91	1.16	9.2		1.77	11.4
			123.2	5.91	1.16	9.2		1.77	11.4
			123.8	5.91	1.16	9.2		1.77	11.4
			124.4	5.91	1.16	9.2		1.77	11.4
			125.0	5.91	1.16	9.2		1.77	11.4
			125.6	5.91	1.16	9.2		1.77	11.4
			126.1	5.91	1.16	9.2		1.77	11.4
			126.7	5.91	1.16	9.2		1.77	11.4
			127.3	5.91	1.16	9.2		1.77	11.4
			127.9	5.91	1.16	9.2		1.77	11.4
			128.5	5.91	1.16	9.2		1.77	11.4
			129.1	5.91	1.16	9.2		1.77	11.4
			129.6	5.91	1.16	9.2		1.77	11.4
			130.2	5.91	1.16	9.2		1.77	11.4
			130.8	5.91	1.16	9.2		1.77	11.4
			131.4	5.91	1.16	9.2		1.77	11.4
			132.0	5.91	1.16	9.2		1.77	11.4
			132.6	5.91	1.16	9.2		1.77	11.4
			133.2	5.91	1.16	9.2		1.77	11.4
			133.8	5.91	1.16	9.2		1.77	11.4
			134.4	5.91	1.16	9.2		1.77	11.4
			135.0	5.91	1.16	9.2		1.77	11.4
			135.6	5.91	1.16	9.2		1.77	11.4
			136.1	5.91	1.16	9.2		1.77	11.4
			136.7	5.91	1.16	9.2		1.77	11.4
			137.3	5.91	1.16	9.2		1.77	11.4
			137.9	5.91	1.16	9.2		1.77	11.4
			138.5	5.91	1.16	9.2		1.77	11.4
			139.1	5.91	1.16	9.2		1.77	11.4
			139.6	5.91	1.16	9.2		1.77	11.4
			140.2	5.91	1.16	9.2		1.77	11.4
			140.8	5.91	1.16	9.2		1.77	11.4
			141.4	5.91	1.16	9.2		1.77	11.4
			142.0	5.91	1.16	9.2		1.77	11.4
			142.6	5.91	1.16	9.2		1.77	11.4
			143.2	5.91	1.16	9.2		1.77	11.4
			143.8	5.91	1.16	9.2		1.77	11.4
			144.4	5.91	1.16	9.2		1.77	11.4
			145.0	5.91	1.16	9.2		1.77	11.4
			145.6	5.91	1.16	9.2		1.77	11.4
			146.1	5.91	1.16	9.2		1.77	11.4
			146.7	5.91	1.16	9.2		1.77	11.4
			147.3	5.91	1.16	9.2		1.77	11.4
			147.9	5.91	1.16	9.2		1.77	11.4
			148.5	5.91	1.16	9.2		1.77	11.4
			149.1	5.91	1.16	9.2		1.77	11.4
			149.7	5.91	1.16	9.2		1.77	11.4
			150.3	5.91	1.16	9.2		1.77	11.4
			150.8	5.91	1.16	9.2		1.77	11.4
			151.4	5.91	1.16	9.2		1.77	11.4
			152.0	5.91	1.16	9.2		1.77	11.4
			152.6	5.91	1.16	9.2		1.77	11.4
			153.2	5.91	1.16	9.2		1.77	11.4
			153.8	5.91	1.16	9.2		1.77	11.4
			154.4	5.91	1.16	9.2		1.77	11.4
			155.0	5.91	1.16	9.2		1.77	11.4
			155.6	5.91	1.16	9.2		1.77	11.4
			156.1	5.91	1.16	9.2		1.77	11.4
			156.7	5.91	1.16	9.2		1.77	11.4
			157.3	5.91	1.16	9.2		1.77	11.4
			157.9	5.91	1.16	9.2		1.77	11.4
			158.5	5.91	1.16	9.2		1.77	11.4
			159.1	5.91	1.16	9.2		1.77	11.4
			159.6	5.91	1.16	9.2		1.77	11.4
			160.2	5.91	1.16	9.2		1.77	11.4
			160.8	5.91	1.16	9.2		1.77	11.4
			161.4	5.91	1.16	9.2		1.77	11.4
			162.0	5.91	1.16	9.2		1.77	11.4
			162.6	5.91	1.16	9.2		1.77	11.4
			163.2	5.91	1.16	9.2		1.77	11.4
			163.8	5.91	1.16	9.2		1.77	11.4
			164.4	5.91	1.16	9.2		1.77	11.4
			165.0	5.91	1.16	9.2		1.77	11.4
			165.6	5.91	1.16	9.2		1.77	11.4
			166.1	5.91	1.16	9.2		1.77	11.4
			166.7	5.91	1.16	9.2		1.77	11.4
			167.3	5.91	1.16	9.2		1.77	11.4
			167.9	5.91	1.16	9.2		1.77	11.4
			168.5	5.91	1.16	9.2		1.77	11.4
			169.1	5.91	1.16	9.2		1.77	11.4
			169.6	5.91	1.16	9.2		1.77	11.4
			170.2	5.91	1.16	9.2		1.77	11.4
			170.8	5.91	1.16	9.2		1.77	11.4
			171.4	5.91	1.16	9.2		1.77	11.4
			172.0	5.91	1.16	9.2		1.77	11.4
			172.6	5.91	1.16	9.2		1.77	11.4
			173.2	5.91	1.16	9.2		1.77	11.4
			173.8	5.91	1.16	9.2		1.77	11.4
			174.4	5.91	1.16	9.2		1.77	11.4
			175.0	5.91	1.16	9.2		1.77	11.4
			175.6	5.91	1.16	9.2		1.77	11.4
			176.1	5.91	1.16	9.2		1.77	11.4
			176.7	5.91	1.16	9.2		1.77	11.4
			177.3	5.91	1.16	9.2		1.77	11.4
			177.9	5.91	1.16	9.2		1.77	11.4
			178.5	5.91	1.16	9.2		1.77	11.4
			179.1	5.91	1.16	9.2		1.77	11.4
			179.6	5.91	1.16	9.2		1.77	11.4
			180.2	5.91	1.16	9.2		1.77	11.4
			180.8	5.91	1.16	9.2		1.77	11.4
			181.4	5.91	1.16	9.2		1.77	11.4
			182.0	5.91	1.16	9.2		1.77	11.4
			182.6	5.91	1.16	9.2		1.77	11.4
			183.2	5.91	1.16	9.2		1.77	11.4
			183.8	5.91	1.16	9.2		1.77	11.4
			184.4	5.91	1.16	9.2		1.77	11.4
			185.0	5.91	1.16	9.2		1.77	11.4
			185.6	5.91	1.16	9.2		1.77	11.4
			186.1	5.91	1.16	9.2		1.77	11.4
			186.7	5.91	1.16	9.2		1.77	11.4
			187.3	5.91	1.16	9.2		1.77	11.4
			187.9	5.91	1.16	9.2		1.77	11.4
			188.5	5.91	1.16	9.2		1.77	11.4
			189.1	5.91	1.16	9.2		1.77	11.4
			189.6	5.91	1.16	9.2		1.77	11.4
			190.2	5.91	1.16	9.2		1.77	11.4
			190.8	5.91	1.16	9.2		1.77	11.4
			191.4	5.91	1.16	9.2		1.77	11.4
			192.0	5.91	1.16	9.2		1.77	11.4
			192.6	5.91	1.16	9.2		1.77	11.4
			193.2	5.91	1.16	9.2		1.77	11.4
			193.8	5.91	1.16	9.2		1.77	11.4
			194.4	5.91	1.16	9.2		1.77	11.4
			195.0	5.91	1.16	9.2		1.77	11.4
			195.6	5.91	1.16	9.2		1.77	11.4
			196.1	5.91	1.16	9.2		1.77	11.4
			196.7	5.91	1.16	9.2		1.77	11.4
			197.3	5.91	1.16	9.2		1.77	11.4
			197.9	5.91	1.16	9.2		1.77	11.4
			198.5	5.91	1.16	9.2		1.77	11.4
			199.1	5.91	1.16	9.2		1.77	11.4
			199.6	5.91	1.16	9.2		1.77	11.4
			200.2	5.91	1.16	9.2		1.77	11.4
			200.8	5.91	1.16	9.2		1.77	11.4
			201.4	5.91	1.16	9.2		1.77	11.4
			202.0	5.91	1.16	9.2		1.77	11.4
			202.6	5.91	1.16	9.2		1.77	11.4
			203.2	5.91	1.16	9.2		1.77	11.4
			203.8	5.91	1.16	9.2		1.77	11.4
			204.4	5.91	1.16	9.2		1.77	11.4
			205.0	5.91	1.16	9.2		1.77	11.4
			205.6	5.91	1.16	9.2		1.77	11.4
			206.1	5.91	1.16	9.2		1.77	11.4
			206.7	5.91	1.16	9.2		1.77	11.4
			207.3	5.91	1.16	9.2		1.77	11.4
			207.9	5.91	1.16	9.2		1.77	11.4
			208.5	5.91	1.16	9.2		1.77	11.4
			209.1	5.91	1.16	9.2		1.77	11.4
			209.6	5.91	1.16	9.2		1.77	11.4
			210.2	5.91	1.16	9.2		1.77	11.

R/V ACANIA 30 APR / 1 MAY 79 CHEMICAL MUSCSCALE (CRUIS- VI)									
TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	H03 UM	P04 PK	NOTE: PATIO H03/P04	ATP NG/L	CHL A MG/W3	TEMP DEG C
1300	36 6.5	121 59.1	343.7	20.14	1.84	10.9	731.0	1.33	10.0
			344.2	21.25	1.91	11.1		1.14	9.5
			344.8	22.37	1.99	11.2		0.76	9.5
			345.4	22.55	2.01	11.3		0.63	9.5
			346.0	22.74	2.01	11.3		0.47	9.8
			347.2	22.92	2.00	11.2		0.50	9.8
			348.3	22.93	2.00	11.2		0.51	9.8
			349.9	22.95	2.02	11.2		0.71	9.6
			350.1	22.55	2.03	11.1		0.72	9.5
			350.7	22.55	2.04	11.1		0.77	9.5
			351.2	22.92	2.05	11.2		0.54	9.4
			351.8	22.92	2.04	11.2		0.54	9.4
1400	36 13.6	121 52.7	352.4	22.92	2.05	11.2	762.0	0.70	9.4
			353.0	22.92	2.05	11.2		0.68	9.4
			353.6	22.92	2.06	11.2		0.66	9.4
			354.2	22.92	2.06	11.2		0.66	9.4
			354.7	22.92	2.06	11.2		0.66	9.4
			355.3	22.92	2.06	11.2		0.63	9.3
			355.9	22.92	2.06	11.2		0.65	9.3
			356.5	22.92	2.06	11.2		0.51	9.7
			357.1	21.81	1.93	11.0		0.55	9.7
			358.2	21.62	1.97	11.0		0.57	9.7
			358.8	20.33	1.91	10.6		0.46	9.7
			359.4	20.33	1.91	10.6		0.54	9.9
1500	36 13.6	121 52.7	360.0	20.33	1.91	10.5	757.0	0.31	10.0
			360.6	20.14	1.91	10.5		0.31	10.0
			361.2	20.33	1.91	10.5		0.39	10.0
			361.7	20.51	1.93	10.7		519.0	10.0
			362.3	20.70	1.94	10.6			10.0
			362.9	21.07	1.95	10.8			9.8
			363.5	21.25	1.91	10.8			9.8
			364.1	21.44	1.96	10.9			9.8
			364.7	21.44	1.96	10.9			9.7
			365.2	21.45	1.97	10.8			9.7
			365.8	20.88	1.97	10.5			9.7
			366.4	20.70	1.97	10.5			9.7
							673.0	0.44	

CHEMICAL AND PHYSICAL CHARACTERISTICS - VII

15-8

**121 57.1**

.....  
.....  
.....  
.....

R/V ACANIA

13 JUNE 79

CHEMICAL MESOSCALE CRUISE VIII

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	MC3 C/M	PO4 C/M	NUTR. RATIO N/P34	ATP NG/L	CHL C/M	TEMP C
2200	36 20.4	121 57.1	24.8	10.91	1.09	10.0	1838.0	1.97	11.7
			25.3	11.09	1.13	9.8		1.97	11.7
			25.9	11.17	1.20	9.8		1.84	11.7
			26.4	11.70	1.21	9.7		1.80	11.6
			27.0	11.63	1.23	9.5	917.0	1.76	11.6
			28.1					1.71	11.5
			28.7					1.61	11.5
			29.2					1.57	11.4
			29.9	11.05	1.18	9.5	1276.0	1.55	11.4
			30.4	11.23	1.02	11.1		1.53	11.4
			31.0	11.31	1.04	11.1		1.51	11.4
			31.6	11.45	1.03	11.2		1.49	11.4
			32.2	11.63	1.05	10.1	916.0	1.48	11.4
			32.8	11.78	1.10	10.8		1.44	11.4
			33.4	12.05	1.29	9.3		1.41	11.3
2300	36 11.0	121 52.6	33.9	12.09	1.32	9.2		1.37	11.3
			34.5	11.92	1.29	9.1	1276.0	1.36	11.4
			35.1	11.74	1.27	8.5		1.34	11.4
			35.7	11.40	1.27	8.6		1.32	11.4
			36.3	11.41	1.27	8.6		1.29	11.5
			36.9	10.87	1.24	8.5	1175.0	1.27	11.6
			37.5	10.66	1.24	8.4		1.25	11.6
			38.1	10.51	1.22			1.23	11.6
			38.7	10.19				1.21	11.6
			39.3					1.18	11.5
			39.9					1.15	11.5
			40.5					1.13	11.5
			41.1					1.11	11.5
			41.7					1.09	11.5
			42.3					1.07	11.5
2300	36 11.0	121 52.6	42.9	8.53	1.24	6.2		1.05	11.5
			43.5	8.60	1.20	7.2	1897.0	1.03	11.5
			44.1	8.78	1.18	7.3		1.01	11.5
			44.7	8.35	1.15	7.6		1.00	11.5
			45.3	8.13	1.10	7.7		0.99	11.5
			45.9	7.98	1.06	8.0		0.97	11.5
			46.5	7.62	1.00	8.0		0.95	11.5
			47.1	7.30	0.95	8.1	1761.0	0.93	11.5
			47.7	7.48	0.90	8.2		0.91	11.5
			48.3	7.26	0.90	8.1		0.88	11.5
			48.9	6.50	0.88	7.8		0.86	11.5
			49.5	5.53	0.79	7.5	2057.0	0.84	11.5
			50.1	4.30	0.65	6.6		0.81	11.5
			50.7	2.87	0.52	5.7		0.78	11.5
			51.3	2.97				0.76	11.5

R/V ACANIA

13 JUNE 79

CHEMICAL MESOSCALE (CRUISE VII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NC3 UN	FM4 UN	NOTE: RATIO NC3:FM4	ATP NG/L	CM1 NG/M3	TEMP C
2400	36	8.4	121 55.4	2.06	0.46	4.5	1857.0	11.31	13.2
			50.5	1.77	0.43	4.1		8.75	13.2
			50.7	1.34	0.39	3.4		4.26	13.4
			51.1	0.58	0.36	2.7		2.13	13.5
			51.4	0.87	0.35	2.5		5.57	13.6
			51.6	0.76	0.33	2.3		10.24	13.6
			52.0	0.55	0.33	2.0		14.72	13.5
			52.4	0.66	0.37	1.8	1977.0	15.36	13.5
			52.8	0.76	0.35	2.2			13.4
			53.2	0.52	0.41	1.3			13.4
			53.6	0.37	0.45	0.8			13.4
			54.0	0.32	0.45	0.7			13.4
			54.4	0.33	0.51	0.6			13.4
			54.8	0.33	0.56	0.6			13.4
			55.2	0.30	0.56	0.5			13.4
			55.6	0.35	0.62	0.6	1851.0		13.4
			56.0	0.48	0.63	0.8			13.4
			56.4	0.55	0.63	0.9			13.4
			56.8	0.47	0.70	1.5			13.4
			57.2	0.37	0.65	1.8	1421.0		13.4
			57.6	0.51	0.61	0.8			13.4
			58.0	0.33	0.64	0.5			13.4
			58.4	0.41					13.4
			58.8	0.58					13.4
			59.2	0.53			1851.0		13.4
			59.6	0.59					13.4
			60.0	0.44					13.4
			60.4	0.60					13.4
			60.8	0.61					13.4
			61.2	0.57					13.4
			61.6	0.62					13.4
			62.0	0.59					13.4
			62.4	0.62					13.4
			62.8	0.62					13.4
			63.2	0.62					13.4
			63.6	0.62					13.4
			64.0	0.62					13.4
			64.4	0.62					13.4
			64.8	0.62					13.4
			65.2	0.62					13.4
			65.6	0.62					13.4
			66.0	0.62					13.4
			66.4	0.62					13.4
			66.8	0.62					13.4
			67.2	0.62					13.4
			67.6	0.62					13.4
			68.0	0.62					13.4
			68.4	0.62					13.4
			68.8	0.62					13.4
			69.2	0.62					13.4
			69.6	0.62					13.4
			70.0	0.62					13.4
			70.4	0.62					13.4
			70.8	0.62					13.4
			71.2	0.62					13.4
			71.6	0.62					13.4
			72.0	0.62					13.4
			72.4	0.62					13.4
			72.8	0.62					13.4
			73.2	0.62					13.4
			73.6	0.62					13.4
			74.0	0.62					13.4
			74.4	0.62					13.4
			74.8	0.62					13.4
			75.2	0.62					13.4
			75.6	0.62					13.4
			76.0	0.62					13.4
			76.4	0.62					13.4
			76.8	0.62					13.4
			77.2	0.62					13.4
			77.6	0.62					13.4
			78.0	0.62					13.4
			78.4	0.62					13.4
			78.8	0.62					13.4
			79.2	0.62					13.4
			79.6	0.62					13.4
			80.0	0.62					13.4
			80.4	0.62					13.4
			80.8	0.62					13.4
			81.2	0.62					13.4
			81.6	0.62					13.4
			82.0	0.62					13.4
			82.4	0.62					13.4
			82.8	0.62					13.4
			83.2	0.62					13.4
			83.6	0.62					13.4
			84.0	0.62					13.4
			84.4	0.62					13.4
			84.8	0.62					13.4
			85.2	0.62					13.4
			85.6	0.62					13.4
			86.0	0.62					13.4
			86.4	0.62					13.4
			86.8	0.62					13.4
			87.2	0.62					13.4
			87.6	0.62					13.4
			88.0	0.62					13.4
			88.4	0.62					13.4
			88.8	0.62					13.4
			89.2	0.62					13.4
			89.6	0.62					13.4
			90.0	0.62					13.4
			90.4	0.62					13.4
			90.8	0.62					13.4
			91.2	0.62					13.4
			91.6	0.62					13.4
			92.0	0.62					13.4
			92.4	0.62					13.4
			92.8	0.62					13.4
			93.2	0.62					13.4
			93.6	0.62					13.4
			94.0	0.62					13.4
			94.4	0.62					13.4
			94.8	0.62					13.4
			95.2	0.62					13.4
			95.6	0.62					13.4
			96.0	0.62					13.4
			96.4	0.62					13.4
			96.8	0.62					13.4
			97.2	0.62					13.4
			97.6	0.62					13.4
			98.0	0.62					13.4
			98.4	0.62					13.4
			98.8	0.62					13.4
			99.2	0.62					13.4
			99.6	0.62					13.4
			100.0	0.62					13.4





R/V ACANIA

13 JUNE 79

CHEMICAL MESOSCAL-ECRHS- VII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PN4 UM	MUTP-DATIN NO3/PN4	STD MG/L	CHL 2 MG/M3	TEMP DEG C
0248	36	21.8	122	1.9					
			80.2	11.49	1.44	10.3			11.7
			80.4	11.23	1.00	11.2			11.7
			80.6	11.23	0.98	9.5			11.7
			80.8	11.52	1.17	10.1			11.7
			81.1	11.13	1.10				11.7
			81.3						11.7
			81.5						11.7
			81.7						11.8
			81.9						11.8
			82.4						11.8
			82.9						11.8
			83.4						11.6
			83.9						11.7
			84.5						11.7
			85.0						11.7

CHEMICAL ABSTRACTS (CRUIS- VIII)

CHIL  
PDS/42

THE  
1940

122 2.1

[illegible]

R/V ACARIA

07 AUG / 09 AUG 79

CHEMICAL MESOSCALICRUISE VIII

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	NO4 UM	NUTR-RATIO NO3/PO4	ATP NG/L	CHL A MG/M3	TEMP C
0300	36 26.7	122 9.9	32.3	1.41	0.0	0.0		5.05	13.7
			33.0	1.27	0.0	0.0		5.20	13.7
			33.6	1.73	1.04	1.7		4.65	13.7
			34.2					3.96	13.7
			35.5					3.36	13.7
			36.1	2.52	0.93	2.7		3.36	13.7
			36.4	1.73	0.93	1.9		3.66	13.7
			37.4	1.27	1.05	1.2		4.14	13.7
			38.0	0.55	1.21	0.5		4.13	13.7
			39.7	0.20	1.19	0.2		3.96	13.7
			39.3	1.27	1.25	1.0		3.97	13.7
			40.6	1.62	1.22	0.8		4.02	13.7
			41.2	1.80	1.11	1.6		4.02	13.7
			41.8					4.16	13.7
			42.2					4.39	13.7
0400	36 19.2	122 17.6	43.1	8.52	0.92	9.3		4.06	13.7
			43.7	8.05	0.92	8.7		4.15	13.7
			44.4	8.59	0.95	10.7		4.13	13.7
			45.0	10.20	1.01	10.7		5.01	13.7
			45.6	10.66	1.15	16.2		5.04	13.7
			46.3	10.91	1.23	17.0		5.04	13.7
			47.6	20.91	1.27	0.0		4.02	13.7
			48.2	0.0		0.0		3.94	13.7
			48.8					4.47	13.7
			49.5					4.41	13.7
			50.1	18.20	0.0	0.0		3.60	13.7
			51.7	18.23	0.0	0.0		3.67	13.7
			52.0	18.59	0.0	0.0		3.69	13.7
			53.3					3.69	13.7
			54.5					2.39	13.7
			55.4					2.31	13.7
			55.8					2.01	13.7
			56.4					1.70	13.7
			57.1					1.71	13.7
			57.7					1.63	13.7
			58.0					1.67	13.7
			59.0					1.47	13.7
			60.2					1.41	13.7
			61.5					1.27	13.7
			62.1					1.13	13.7
			62.8					1.01	13.7
			63.4					1.05	13.7

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE K UM	NO3 UM	PO4 UM	NUTR. RATIO NO3/PO4	ATP NG/L
			64.0				

Distance	UM	PO4	NUTR. RATIO	ATP	CHL. a	TEW
km		UM	NU3/PO4	NG/L	mg/L	g/g
64.0					0.91	13.3
64.7					0.84	13.3
65.3					0.445	13.2
65.9					0.480	13.5
66.6					0.719	13.6
67.2					0.75	13.7
67.8					0.75	13.9
68.3					0.80	14.0
68.9					0.79	14.0
69.5					0.74	14.1
70.0					0.74	14.1
70.6					0.82	14.1
71.2					0.83	14.1
71.7					0.98	14.0
72.3					0.97	14.0
72.9					0.53	14.0
73.4					0.95	14.1
74.0					0.98	14.1
74.6					1.02	14.1
75.1					1.03	14.1
75.7					1.05	14.1
76.2					1.05	14.1
76.8					1.05	14.1
77.4					1.04	14.1
77.9					0.97	14.1
78.5					1.04	14.1
79.1					0.99	14.0
79.6					0.97	14.0
80.2					0.97	14.0
80.8					0.95	14.0
81.3					0.95	14.1
81.9					0.95	14.1
82.5					0.93	14.0
83.0					0.97	14.0
83.6					0.97	14.0
84.2					0.97	14.0
84.7					0.97	14.0
85.3					0.97	14.0
85.9					0.97	14.0
86.4					0.97	14.0
87.0					0.97	14.0
87.6					0.96	14.0
88.1					0.97	14.0
89.3					0.97	14.0
89.9					0.90	14.0
90.4					0.90	14.0
90.9					0.93	14.0
91.5					0.93	14.0
92.1					0.55	14.0

R/V ACANIA

07 AUG / 09 AUG 79

CHEMICAL MESOSCALE (CRUISE VIII)

TIME  
GMT  
0600LATITUDE  
NORTH  
36 2.1LONGITUDE  
WEST  
122 32.0DISTANCE  
KMPD3  
UMPD4  
UMNUCLEO-PATIO  
NO3/PD4ATP  
NG/LCHL A  
MG/M3TEMP  
C

0700

35 54.0

122 37.7

92.6	8.15	0.97	8.4	0.96	13.9
93.8	7.78	0.77	10.1	0.96	13.9
94.3				0.97	13.9
94.9				0.97	13.9
95.5				0.97	13.9
96.0	7.74	0.92	8.4	0.97	13.9
96.6	7.67	0.93	8.4	0.97	13.9
97.2	8.17	0.97	8.4	0.97	13.9
97.7	8.16	1.21	6.7	0.97	13.9
98.3	8.28	2.07	4.0	0.97	13.9
99.9				0.97	13.9
100.6				0.97	13.9
101.1				0.97	13.9
101.7				0.97	13.9
102.3				0.97	13.9
102.8				0.97	13.9
103.4				0.97	13.9
103.9				0.97	13.9
104.5				0.97	13.9
105.1				0.97	13.9
105.6				0.97	13.9
106.2				0.97	13.9
106.8				0.97	13.9
107.3				0.97	13.9
107.9				0.97	13.9
108.5				0.97	13.9
109.0				0.97	13.9
109.6				0.97	13.9
110.2				0.97	13.9
110.7				0.97	13.9
111.3				0.97	13.9
111.9				0.97	13.9
112.4				0.97	13.9
113.0				0.97	13.9
113.6				0.97	13.9
114.1				0.97	13.9
114.7				0.97	13.9
115.3				0.97	13.9
115.8				0.97	13.9
116.4				0.97	13.9
117.0				0.97	13.9
117.5				0.97	13.9
118.1				0.97	13.9
118.6				0.97	13.9
119.2				0.97	13.9
119.8				0.97	13.9
120.3				0.97	13.9

R/V ALANIA

07 AUG / 09 AUG 79

CHEMICAL MESOSCALE(CRUISE VITI)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	N <sub>3</sub> UM	PO <sub>4</sub> UM	NUTR. RATIO N <sub>3</sub> /PO <sub>4</sub>	ATP NG/L	CHL. A MG/93	TEMP C
			149.6	7.52	0.0	0.0		0.52	14.5
			150.1	8.73	0.0	0.0		0.45	14.5
			150.6	8.70	0.0	0.0		0.43	14.5
			151.1	8.25	0.0	0.0		0.74	14.5
			151.6	8.77	0.0	0.0		0.78	14.5
			152.0	8.96	0.0	0.0		0.66	14.5
			152.5	8.70	0.0	0.0		0.69	14.5
			153.0	7.66	0.0	0.0		0.72	14.5
			153.5	7.55	0.0	0.0		0.73	14.5
			154.0	7.14	0.0	0.0		0.74	14.5
			154.5	7.14	0.0	0.0		0.74	14.5
			155.0	7.09	0.0	0.0		0.68	14.5
			155.5	7.27	0.0	0.0		0.69	14.5
			156.0	7.32	0.0	0.0		0.67	14.5
			156.4					0.67	14.5
			157.0					0.66	14.5
			157.4					0.63	14.5
			157.9					0.62	14.5
			158.4					0.62	14.5
			159.0					0.62	14.5
			160.0					0.61	14.5
			160.3					0.59	14.5
			161.0					0.57	14.5
			161.3					0.55	14.5
			162.0					0.55	14.5
			162.3					0.52	14.5
			162.8					0.52	14.5
			163.0					0.50	14.5
			163.3					0.50	14.5
			164.0					0.47	14.5
			164.3					0.47	14.5
			165.0					0.44	14.5
			165.3					0.44	14.5
			165.9					0.43	14.5
			166.3					0.43	14.5
			167.0					0.43	14.5
			167.4					0.44	14.5
			168.0					0.43	14.5
			168.4					0.42	14.5
			169.0					0.42	14.5
			169.4					0.45	14.5
			170.0					0.45	14.5
			171.0					0.43	14.5
			171.5					0.43	14.5
			172.0					0.44	14.5
			172.5					0.44	14.5
			173.0					0.48	14.5
			173.5					0.48	14.5
			174.0					0.48	14.5

1000 35 29.1 122 57.1

R/V ACANIA 07 AUG / 09 AUG 79 CHEMICAL MESOSCALTICRUISE VIII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NC3 UM	PC4 UP	NU/P-RATIO NU3/PC4	ATP NG/L	CHL A MG/L	TEMP DEG C
0800	35 42.0	122 44.8	120.9	3.04	0.95	3.2		1.09	15.3
			121.0	3.17	0.89	3.6		1.04	15.3
			122.6	3.86	0.89	3.6		1.04	15.3
			123.7	3.07	0.91	3.4		0.97	15.3
			124.4	3.16	0.89	3.5		0.96	15.3
			125.4	3.41	0.91	3.7		0.91	15.3
			126.0	3.54	0.93	3.8		0.88	15.3
			127.6					0.88	15.3
			127.8					0.84	15.3
			129.0					0.76	15.3
			130.0					0.71	15.3
			130.9					0.69	15.3
			131.5					0.70	15.3
			132.1					0.72	15.3
			133.3					0.70	15.3
			133.9					0.68	15.3
			134.4					0.65	15.3
			135.1					0.62	15.3
			136.0					0.63	15.3
			137.5					0.65	15.3
			138.1					0.66	15.3
			139.3					0.66	15.3
			140.5					0.69	15.3
			141.1					0.70	15.3
			141.3					0.77	15.3
			142.3					0.77	15.3
			143.5					0.81	15.3
			144.1					0.85	15.3
			144.7					0.85	15.3
			145.2					0.86	15.3
			146.2					0.84	15.3
			146.7					0.85	15.3
			147.2					0.84	15.3
			147.6					0.83	15.3
			148.1					0.82	15.3
			149.1					0.59	14.7
0900	35 36.4	122 49.9	3.66	0.94	3.9			3.68	15.3
			6.13	0.90	6.9			6.85	15.3
			6.41	0.92	6.7			6.85	15.3
			6.70	0.95	6.7			6.85	15.3
			6.85	0.98	6.8			6.85	15.3
			7.14	0.91	7.4			6.85	15.3
			7.40	0.0	0.0			0.83	14.7
			7.89	0.0	0.0			0.83	14.7
			8.12	0.0	0.0			0.82	14.7
			7.92	0.0	0.0			0.59	14.7



TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE K	NO3 UM	PO4 UM	NUTR. RATIO NO3/PO4	ATP NG/L
1100	35 20.6	123 2.7	174.6	175.1			

1200	35	12.1	123	8.1
------	----	------	-----	-----

Distance	NO3	PO4	NUTR. RATIO NO3/PO4	ATP NG/L	CHL A MG/L	TPP MG/L
174.6	UM	UM			0.49	15.3
175.1					0.50	15.3
175.4					0.50	15.3
176.1					0.50	15.3
175.6					0.50	15.3
177.1					0.48	15.3
177.7					0.49	15.2
174.2					0.50	15.2
178.6					0.50	15.2
179.1					0.49	15.2
179.6					0.48	15.1
180.1					0.49	15.1
180.5					0.37	15.1
181.0					0.32	15.0
181.5					0.32	14.9
181.9					0.32	14.9
182.4					0.32	14.9
183.2					0.32	14.8
183.6					0.34	14.8
184.3					0.37	14.8
184.8					0.45	14.8
185.3					0.47	14.5
185.7					0.49	14.5
186.4					0.50	14.4
187.0					0.51	14.4
187.6					0.51	14.4
188.3					0.46	14.4
189.9					0.46	14.4
190.2					0.40	14.4
190.8					0.39	14.6
191.4					0.41	14.7
192.1					0.47	14.7
192.7					0.63	14.8
193.3					0.68	14.8
194.0					0.68	14.8
194.6					0.69	14.8
195.2					0.69	14.8
195.9					0.73	14.9
196.5					0.73	14.9
197.1					0.74	14.9
197.8					0.74	14.9
198.4					0.73	15.0
199.0					0.74	15.0
199.7					0.74	15.0
200.3					0.75	15.0
200.9					0.75	15.0
201.6					0.75	15.1
202.2					0.77	15.1

R/V ACARUA

07 AUG / 09 AUG 79

CHEMICAL MESOSCALE (CRUISE VIII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	M03 UM	PD4 UM	NUTR. RATIO NO3/PO4	ATP NG/L	CHL. A PG/ML	TEMP DEG. C
1300	35	3.7	123 13.7	202.8	11.11	0.71	14.2	0.19	15.1
				203.5	11.25	0.74	14.4	0.17	15.1
				204.1	11.02	0.74	14.9	0.17	15.1
				205.4	10.74	0.73	15.1	0.17	15.2
				206.0	10.45	0.69	15.1	0.17	15.2
				207.3	10.59	0.68	15.6	0.17	15.1
				207.3	10.05	0.69	15.6	0.17	15.1
				208.5	9.88	0.68	14.5	0.17	15.1
				209.2	9.73	0.67	14.5	0.17	15.1
				209.3	9.65	0.76	14.5	0.17	15.1
				210.5	9.71	0.67	14.1	0.17	15.1
				211.1	9.76	0.67	14.1	0.17	15.1
1400	34	55.2	123 19.4	212.4	9.39	0.63	13.8	0.17	15.1
				213.0	9.88	0.70	14.1	0.17	15.2
				214.3	10.02	0.70	14.3	0.17	15.2
				214.9	9.59	0.69	13.9	0.17	15.2
				215.5	10.02	0.69	14.5	0.17	15.2
				216.2	10.16	0.68	14.5	0.17	15.2
				216.8	10.02	0.67	15.0	0.17	15.3
				217.4	9.36	0.69	13.3	0.17	15.3
				218.1	9.96	0.67	14.9	0.17	15.3
				219.7	9.96	0.67	14.6	0.17	15.3
				220.0	9.59	0.68	14.6	0.17	15.3
				220.6	9.59	0.68	14.5	0.17	15.3
1500	34	55.2	123 19.4	221.2	10.16	0.69	14.5	0.17	15.3
				222.5	10.02	0.67	14.5	0.17	15.3
				223.1	9.59	0.67	14.5	0.17	15.3
				223.4	9.59	0.67	14.5	0.17	15.3
				224.4	9.59	0.67	14.5	0.17	15.3
				225.0	9.59	0.67	14.5	0.17	15.3
				225.7	9.59	0.67	14.5	0.17	15.3
				226.1	9.59	0.67	14.5	0.17	15.3
				227.0	9.59	0.67	14.5	0.17	15.3
				227.6	9.59	0.67	14.5	0.17	15.3
				228.1	9.59	0.67	14.5	0.17	15.3
				229.2	9.59	0.67	14.5	0.17	15.3

R/V ACANIA

07 AUG / 09 AUG 79

CHEMICAL MESUSCALE CRUISE VIII

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NC3 UM	PC4 UM	NUTR. RATIO NO3/PO4	ATP MG/L	CHL MG/M3	TEMP C
1452	35 47.3	123 24.7	234.5	8.02	0.05	9.4	1.02	1.24	15.5
			235.2	7.87	0.05	9.4	1.02	1.24	15.5
			236.4	7.73	0.05	9.4	1.02	1.24	15.5
			237.1	7.59	0.05	9.4	1.02	1.24	15.5
			238.3	7.44	0.05	9.4	1.02	1.24	15.5
			239.0	7.29	0.05	9.4	1.02	1.24	15.5
			240.2	7.10	0.05	9.4	1.02	1.24	15.5
			241.5	6.96	0.05	9.4	1.02	1.24	15.5
			242.2	6.81	0.05	9.4	1.02	1.24	15.5
			243.8	6.66	0.05	9.4	1.02	1.24	15.5
			244.4	6.53	0.05	9.4	1.02	1.24	15.5
			245.3	6.40	0.05	9.4	1.02	1.24	15.5
			246.0	6.25	0.05	9.4	1.02	1.24	15.5
			247.2	6.10	0.05	9.4	1.02	1.24	15.5
			248.1	5.96	0.05	9.4	1.02	1.24	15.5
			249.5	5.81	0.05	9.4	1.02	1.24	15.5
			250.4	5.67	0.05	9.4	1.02	1.24	15.5
			251.0	5.52	0.05	9.4	1.02	1.24	15.5
			252.4	5.37	0.05	9.4	1.02	1.24	15.5
			253.7	5.22	0.05	9.4	1.02	1.24	15.5
			254.9	5.07	0.05	9.4	1.02	1.24	15.5
			255.4	4.92	0.05	9.4	1.02	1.24	15.5
			256.1	4.77	0.05	9.4	1.02	1.24	15.5
			257.4	4.62	0.05	9.4	1.02	1.24	15.5
			258.9	4.47	0.05	9.4	1.02	1.24	15.5
			259.9	4.32	0.05	9.4	1.02	1.24	15.5
			260.5	4.17	0.05	9.4	1.02	1.24	15.5
			261.2	4.02	0.05	9.4	1.02	1.24	15.5
			261.8	3.87	0.05	9.4	1.02	1.24	15.5
			262.4	3.72	0.05	9.4	1.02	1.24	15.5
			263.1	3.57	0.05	9.4	1.02	1.24	15.5
			263.7	3.42	0.05	9.4	1.02	1.24	15.5
			264.3	3.27	0.05	9.4	1.02	1.24	15.5
			265.0	3.12	0.05	9.4	1.02	1.24	15.5
			265.6	2.97	0.05	9.4	1.02	1.24	15.5
			266.3	2.82	0.05	9.4	1.02	1.24	15.5
			266.9	2.67	0.05	9.4	1.02	1.24	15.5
			267.5	2.52	0.05	9.4	1.02	1.24	15.5
			268.2	2.37	0.05	9.4	1.02	1.24	15.5
			268.8	2.22	0.05	9.4	1.02	1.24	15.5
			269.5	2.07	0.05	9.4	1.02	1.24	15.5
			270.1	1.92	0.05	9.4	1.02	1.24	15.5
			270.8	1.77	0.05	9.4	1.02	1.24	15.5
			271.4	1.62	0.05	9.4	1.02	1.24	15.5
			272.1	1.47	0.05	9.4	1.02	1.24	15.5
			272.7	1.32	0.05	9.4	1.02	1.24	15.5
			273.4	1.17	0.05	9.4	1.02	1.24	15.5
			274.0	1.02	0.05	9.4	1.02	1.24	15.5
			274.7	0.87	0.05	9.4	1.02	1.24	15.5
			275.3	0.72	0.05	9.4	1.02	1.24	15.5
			276.0	0.57	0.05	9.4	1.02	1.24	15.5
			276.6	0.42	0.05	9.4	1.02	1.24	15.5
			277.3	0.27	0.05	9.4	1.02	1.24	15.5
			277.9	0.12	0.05	9.4	1.02	1.24	15.5
			278.6	0.00	0.05	9.4	1.02	1.24	15.5
			279.2	0.00	0.05	9.4	1.02	1.24	15.5
			279.9	0.00	0.05	9.4	1.02	1.24	15.5
			280.5	0.00	0.05	9.4	1.02	1.24	15.5
			281.2	0.00	0.05	9.4	1.02	1.24	15.5
			281.8	0.00	0.05	9.4	1.02	1.24	15.5
			282.4	0.00	0.05	9.4	1.02	1.24	15.5
			283.1	0.00	0.05	9.4	1.02	1.24	15.5
			283.7	0.00	0.05	9.4	1.02	1.24	15.5
			284.3	0.00	0.05	9.4	1.02	1.24	15.5
			285.0	0.00	0.05	9.4	1.02	1.24	15.5
			285.6	0.00	0.05	9.4	1.02	1.24	15.5
			286.3	0.00	0.05	9.4	1.02	1.24	15.5
			286.9	0.00	0.05	9.4	1.02	1.24	15.5
			287.5	0.00	0.05	9.4	1.02	1.24	15.5
			288.2	0.00	0.05	9.4	1.02	1.24	15.5
			288.8	0.00	0.05	9.4	1.02	1.24	15.5
			289.5	0.00	0.05	9.4	1.02	1.24	15.5
			290.1	0.00	0.05	9.4	1.02	1.24	15.5
			290.8	0.00	0.05	9.4	1.02	1.24	15.5
			291.4	0.00	0.05	9.4	1.02	1.24	15.5
			292.1	0.00	0.05	9.4	1.02	1.24	15.5
			292.7	0.00	0.05	9.4	1.02	1.24	15.5
			293.4	0.00	0.05	9.4	1.02	1.24	15.5
			294.0	0.00	0.05	9.4	1.02	1.24	15.5
			294.7	0.00	0.05	9.4	1.02	1.24	15.5
			295.3	0.00	0.05	9.4	1.02	1.24	15.5
			296.0	0.00	0.05	9.4	1.02	1.24	15.5
			296.6	0.00	0.05	9.4	1.02	1.24	15.5
			297.3	0.00	0.05	9.4	1.02	1.24	15.5
			297.9	0.00	0.05	9.4	1.02	1.24	15.5
			298.6	0.00	0.05	9.4	1.02	1.24	15.5
			299.2	0.00	0.05	9.4	1.02	1.24	15.5
			299.9	0.00	0.05	9.4	1.02	1.24	15.5
			300.5	0.00	0.05	9.4	1.02	1.24	15.5
			301.2	0.00	0.05	9.4	1.02	1.24	15.5
			301.8	0.00	0.05	9.4	1.02	1.24	15.5
			302.4	0.00	0.05	9.4	1.02	1.24	15.5
			303.1	0.00	0.05	9.4	1.02	1.24	15.5
			303.7	0.00	0.05	9.4	1.02	1.24	15.5
			304.3	0.00	0.05	9.4	1.02	1.24	15.5
			305.0	0.00	0.05	9.4	1.02	1.24	15.5
			305.6	0.00	0.05	9.4	1.02	1.24	15.5
			306.3	0.00	0.05	9.4	1.02	1.24	15.5
			306.9	0.00	0.05	9.4	1.02	1.24	15.5
			307.5	0.00	0.05	9.4	1.02	1.24	15.5
			308.2	0.00	0.05	9.4	1.02	1.24	15.5
			308.8	0.00	0.05	9.4	1.02	1.24	15.5
			309.5	0.00	0.05	9.4	1.02	1.24	15.5
			310.1	0.00	0.05	9.4	1.02	1.24	15.5
			310.8	0.00	0.05	9.4	1.02	1.24	15.5
			311.4	0.00	0.05	9.4	1.02	1.24	15.5
			312.1	0.00	0.05	9.4	1.02	1.24	15.5
			312.7	0.00	0.05	9.4	1.02	1.24	15.5
			313.4	0.00	0.05	9.4	1.02	1.24	15.5
			314.0	0.00	0.05	9.4	1.02	1.24	15.5
			314.7	0.00	0.05	9.4	1.02	1.24	15.5
			315.3	0.00	0.05	9.4	1.02	1.24	15.5
			316.0	0.00	0.05	9.4	1.02	1.24	15.5
			316.6	0.00	0.05	9.4	1.02	1.24	15.5
			317.3	0.00	0.05	9.4	1.02	1.24	15.5
			317.9	0.00	0.05	9.4	1.02	1.24	15.5
			318.6	0.00	0.05	9.4	1.02	1.24	15.5
			319.2	0.00	0.05	9.4	1.02	1.24	15.5
			319.9	0.00	0.05	9.4	1.02	1.24	15.5
			320.5	0.00	0.05	9.4	1.02	1.24	15.5
			321.2	0.00	0.05	9.4	1.02	1.24	15.5
			321.8	0.00	0.05	9.4	1.02	1.24	15.5
			322.4	0.00	0.05	9.4	1.02	1.24	15.5
			323.1	0.00	0.05	9.4	1.02	1.24	15.5
			323.7	0.00	0.05	9.4	1.02	1.24	15.5
			324.3	0.00	0.05	9.4	1.02	1.24	15.5
			325.0	0.00	0.05	9.4	1.02	1.24	15.5
			325.6	0.00	0.05	9.4	1.02	1.24	15.5
			326.3	0.00	0.05	9.4	1.02	1.24	15.5
			326.9	0.00	0.05	9.4	1.02	1.24	15.5
			327.5	0.00	0.05	9.4	1.02	1.24	15.5
			328.2	0.00	0.05	9.4	1.02	1.24	15.5
			328.8	0.00	0.05	9.4	1.02	1.24	15.5
			329.5	0.00	0.05	9.4	1.02	1.24	15.5
			330.1	0.00	0.05	9.4	1.02	1.24	15.5
			330.8	0.00	0.05	9.4	1.02	1.24	15.5
			331.4	0.00	0.05	9.4	1.02	1.24	15.5
			332.1	0.00	0.05	9.4	1.02	1.24	15.5
			332.7	0.00	0.05	9.4	1.02	1.24	15.5
			333.4	0.00	0.05	9.4	1.02	1.24	15.5
			334.0	0.00	0.05	9.4	1.02	1.24	15.5
			334.7	0.00	0.05	9.4	1.02	1.24	15.5
			335.3	0.00	0.05	9.4	1.02	1.24	15.5
			336.0	0.00	0.05	9.4	1.02	1.24	15.5
			336.6	0.00	0.05	9.4	1.02	1.24	15.5
			337.3	0.00	0.05	9.4	1.02	1.24	15.5
			337.9	0.00	0.05	9.4	1.02	1.24	15.5
			338.6	0.00	0.05	9.4	1.02	1.24	15.5
			339.2	0.00	0.05	9.4	1.02	1.24	15.5
			339.9	0.00	0.05	9.4	1.02	1.24	15.5
			340.5	0.00	0.05	9.4	1.02	1.24	15.5
			341.2	0.00	0.05	9.4	1.02	1.24	15.5
			341.8	0.00	0.05	9.4	1.02	1.24	15.5
			342.4</						

89

R/V ACANIA 07 AUG / 09 AUG 79 CHEMICAL RESIDUALS (CRUISE VIII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 M	PO4 M	NUTR. RATIO NH3/PO4	ATP NG/L	CHL A MG/43	TEMP D.C. C
1805	34 32.8	123 32.3	297.9	0.0	0.25	0.0	788.0	0.17	17.1
			308.6	0.0	0.25	0.0		0.17	17.1
			299.2	0.0	0.23	0.0		0.17	17.1
			300.5	0.0	0.25	0.0		0.17	17.1
			301.7	1.09	0.25	4.4		0.17	17.1
			302.4	0.87	0.25	3.6		0.17	17.1
			303.0	0.89	0.27	3.0		0.17	17.1
			303.6	0.89	0.27	3.3		0.17	17.1
			304.3	0.82	0.27	3.3		0.17	17.1
			305.5	0.68	0.27	3.4		0.17	17.1
			306.2	0.98	0.29	3.5		0.19	17.0
			307.4	0.96	0.25	3.3		0.20	17.0
			308.1			3.8		0.21	17.0
			308.7					0.21	17.0
			309.3	1.13	0.27	4.4	925.0	0.21	17.0
			310.0	0.84	0.26	3.8		0.22	16.9
			311.2	0.93	0.26	3.8		0.26	16.9
			311.5	0.96	0.26	3.7		0.28	16.9
			312.5	0.93	0.25	3.7	1165.0	0.29	16.9
			313.4	0.91	0.25	3.9		0.29	16.9
			314.4	0.83	0.25	3.6		0.33	16.6
			315.7	1.05	0.26	3.7		0.34	16.6
			316.3	1.22	0.32	4.2	1014.0	0.37	16.6
			317.6	1.31	0.32	4.1		0.40	16.6
			318.2	1.10	0.32	3.9		0.41	16.6
			319.5	1.16	0.35	3.9		0.44	16.6
			320.7	1.45	0.35	2.5	3418.0	0.47	16.6
			321.4	0.06	0.45	1.3		0.47	16.6
1855	34 39.4	123 36.0	322.0					0.47	16.6
			322.6					0.47	16.6
1900	34 39.9	123 36.4	323.3					0.43	15.6
			323.9					0.43	15.6
			324.5					0.43	15.6
			325.2					0.43	15.6
			326.4					0.43	15.6
			327.1				1256.0	0.50	15.7
			328.4					0.86	15.6
			329.0					0.86	15.6

R/V ACANIA		07 AUG / 09 AUG 79		CHEMICAL MESOSCALE CRUISE VIII					
TIME GMT	LATITUDE NCRTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR. RATIO NO3/PO4	ATP NG/L	CHL A MC/43	TIME DIG C
1931	34 43.9	123 39.4	329.6	0.0	0.53	0.0	1082.0	0.47	15.6
			330.0	0.0	0.49	0.0		0.47	15.6
			331.5	0.0	0.54	0.0		0.37	17.0
			332.2	0.0	0.49	0.0		0.36	16.8
			333.4	0.0	0.49	0.0		0.36	16.8
			334.1	0.0	0.49	0.0		0.36	16.8
			335.3	0.0	0.49	0.0		0.36	16.8
			336.0	0.0	0.49	0.0		0.36	16.8
			337.6	0.0	0.49	0.0		0.36	16.8
			338.5	0.0	0.49	0.0		0.36	16.8
			339.1	0.0	0.49	0.0		0.36	16.8
			340.4	0.0	0.49	0.0		0.36	16.8
			341.7	0.0	0.49	0.0		0.36	16.8
			342.9	0.0	0.49	0.0		0.36	16.8
2000	34 42.2	123 42.6	343.0	0.0	0.49	0.0		0.36	16.8
			344.2	0.0	0.49	0.0		0.36	16.8
			345.5	0.0	0.49	0.0		0.36	16.8
			346.1	0.0	0.49	0.0		0.36	16.8
			347.4	0.0	0.49	0.0		0.36	16.8
			348.6	0.0	0.49	0.0		0.36	16.8
			349.9	0.0	0.49	0.0		0.36	16.8
			351.2	0.0	0.49	0.0		0.36	16.8
			352.4	0.0	0.49	0.0		0.36	16.8
			353.7	0.0	0.49	0.0		0.36	16.8
			354.3	0.0	0.49	0.0		0.36	16.8
			355.0	0.0	0.49	0.0		0.36	16.8
			356.6	0.0	0.49	0.0		0.36	16.8
			357.9	0.0	0.49	0.0		0.36	16.8
			358.1	0.0	0.49	0.0		0.36	16.8
			359.4	0.0	0.49	0.0		0.36	16.8
			360.7	0.0	0.49	0.0		0.36	16.8

[illegible]

[illegible]



R/V ACANIA

07 AUG / 09 AUG 79

CHEMICAL MESOSCALE (CRUISE VIII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	N/3 UM	P04 UM	NU/P-RATIO NU3/P04	ATP NG/L	CHL A MG/43	TRP DEG C
0100	34	55.4	123 38.1						
			424.7	2.03	0.70	2.9	0.52	0.52	14.0
			425.3	2.16	0.71	3.0	0.52	0.52	14.0
			426.0	2.07	0.71	2.9	0.52	0.52	14.0
			426.6	1.89	0.71	2.7	0.47	0.47	14.0
			427.2	1.21	0.69	1.8	0.45	0.45	13.5
			428.5				0.42	0.42	13.5
			429.1				0.42	0.42	13.5
			429.8				0.42	0.42	13.5
			430.4				0.42	0.42	13.5
			431.0				0.42	0.42	13.5
			431.7				0.42	0.42	13.5
			432.3				0.42	0.42	13.5
			433.0				0.42	0.42	13.5
			433.6				0.42	0.42	13.5
			434.2				0.42	0.42	13.5
			434.8				0.42	0.42	13.5
			435.5				0.42	0.42	13.5
			436.1				0.42	0.42	13.5
			436.7				0.42	0.42	13.5
			437.4				0.42	0.42	13.5
			438.0				0.42	0.42	13.5
			438.6				0.42	0.42	13.5
			439.3				0.42	0.42	13.5
			440.5				0.42	0.42	13.5
			441.2				0.42	0.42	13.5
			441.8				0.42	0.42	13.5
			442.4				0.42	0.42	13.5
			443.1				0.42	0.42	13.5
			443.7				0.42	0.42	13.5
			444.3				0.42	0.42	13.5
			445.0				0.42	0.42	13.5
			445.6				0.42	0.42	13.5
			446.2				0.42	0.42	13.5
			446.9				0.42	0.42	13.5
			447.5				0.42	0.42	13.5
			448.1				0.42	0.42	13.5
			448.8				0.42	0.42	13.5
			449.4				0.42	0.42	13.5
			450.0				0.42	0.42	13.5
			450.7				0.42	0.42	13.5
			451.3				0.42	0.42	13.5
			451.9				0.42	0.42	13.5
			452.6				0.42	0.42	13.5
			453.2				0.42	0.42	13.5
			453.8				0.42	0.42	13.5
			454.5				0.42	0.42	13.5
			455.1				0.42	0.42	13.5
			455.7				0.42	0.42	13.5

TIME GMT 0200	LATITUDE NORTH 35 1.9	LONGITUDE WEST 123 31.7	DISTANCE KM	NC3 HM	PM4 UM	NUTR. RATIO N:P3/PO4	ATP NG/L	PHL 4 M7/93	TEMP DEG C
			456.4	5.65	0.85	6.6		0.75	14.5
			457.0	5.61	0.85	6.6		0.77	15.0
			458.3	5.74	0.83	6.9		0.86	14.9
			459.5					0.83	14.6
			460.4	5.34	0.87	6.1		0.73	14.4
			460.9	5.21	0.85	5.4		0.70	14.5
			461.4	5.39	0.83	6.1		0.73	14.8
			462.7	5.69	0.82	6.5		0.72	14.8
			463.3	5.92	0.85	6.9		0.73	14.5
			464.0	6.14	0.85	7.8		0.73	14.6
			465.0	6.44	0.85	7.9		0.73	14.7
			465.2	6.22	0.85	7.6		0.73	14.7
			465.7	5.17	0.84	6.2		0.73	14.5
			466.5	4.25	0.80	5.3		0.73	14.5
			467.2	2.42	0.74	3.3		0.73	15.0
			467.8					0.73	15.0
			468.4					0.73	15.0
			469.1					0.73	15.0
			469.7					0.73	15.0
			470.3	4.34	1.10	3.9		0.56	15.0
			471.0	3.91	1.09	3.6		0.56	15.0
			471.6	3.69	1.05	3.5		0.56	15.1
			472.2	3.51	0.94	3.6		0.56	15.1
			473.3	3.25	0.97	3.4		0.56	15.2
			474.1	3.03	0.95	3.2		1.01	15.3
			474.8	2.81	0.93	3.0		1.01	15.3
			475.4	2.73	0.93	2.7		1.01	15.4
			476.0	2.60	0.93	2.7		1.01	15.4
			476.7	2.38	0.93	2.6		1.01	15.4
			477.3	1.81	0.92	1.9		1.01	15.4
			477.9	1.29	0.81	1.4		1.01	15.4
			478.6	1.20	0.81	1.5		1.01	15.4
			479.2	1.51	0.79	1.9		1.01	15.4
			479.9	1.72	0.74	2.3		1.01	15.4
			480.5	1.72	0.77	2.2		1.01	15.4
			481.1	1.51	0.75	2.2		1.01	15.4
			481.7					1.01	15.4
			482.4					1.01	15.4
			483.0	1.59	0.77	2.1		1.01	15.4
			483.6	1.72	0.79	2.2		1.01	15.4
			484.3	1.72	0.77	2.2		1.01	15.4
			484.9	1.81	0.77	2.4		1.01	15.4
			485.5	1.94	0.77	2.5		1.01	15.4
			486.2	1.94	0.78	2.5		1.01	15.4
			486.8	2.16		2.8		1.01	15.4
			487.4					1.01	15.4

0300 35 8.6 123 25.8

NO-A089 930

NAVAL POSTGRADUATE SCHOOL MONTEREY CA  
RELATIONSHIPS BETWEEN SEA SURFACE TEMPERATURE AND NUTRIENTS IN -ETC(U)  
MAR 80 J W CONRAD

F/G 17/5


UNCLASSIFIED

NL

2 of 2  
516  
11-80



END  
DATE  
FILMED  
11-80  
DTIC



R/V ACAP'IA 07 AUG / 09 AUG 79 CHEMICAL WETSCALE(CRUISE VII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NC3 UM	PC6 UM	NUTR. RATIO NO3/PO4	ATP NG/L	CHL A PG/43	TEMP C
0415	35 16.9	123 17.3	489.1	2.39	0.79	2.9			15.5
			489.7	0.0	0.80	0.0			15.5
			489.1	0.0	0.81	0.0			15.5
			490.0	0.0	0.81	0.0			15.5
			490.6						15.5
			491.2						15.5
			491.4						15.5
			491.5						15.5
			493.1						15.5
			493.9						15.5
			494.4						15.5
			495.0						15.5
			495.7						15.5
			496.3						15.5
			496.9						15.5
			497.6						15.5
			498.2						15.5
			499.8						15.5
			500.1						15.5
			500.7						15.5
			501.4						15.5
			502.0						15.5
			502.6						15.5
			503.3						15.5
			504.5						15.5
			505.2						15.5
			505.8						15.5
			506.4						15.5
			507.1						15.5
			507.7						15.5
			508.3						15.5
			509.0						15.5
			510.2						15.5
			511.5						15.5
			512.1						15.5
			513.6						15.5
			514.1						15.5
			514.7						15.5
			515.3						15.5
			516.0						15.5
			517.2						15.5
			518.5						15.5
			519.1						15.5
0505.	35 22.7	123 11.7							

CHEMICAL MESOSCALE CRUISES = VIIIIII

CHL A  
MG/M3

0600	35	28.6	123	6.2
------	----	------	-----	-----

[illegible]

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE NM	PD4 UM	NUTR. RATIO NDS/PO4	ATP MG/L	CHL A MG/M3
0700	35 35.0	123 0.6	55.1 ↓	UM			

0800	35 42.6	122 53.9
------	---------	----------

EXP	CHL A	ATP	MUTR. RATIO	PO <sub>4</sub>	NO <sub>3</sub>	DISTANCE
MG/M <sup>3</sup>	NG/L	NO <sub>3</sub> /PO <sub>4</sub>	UM	UM	KM	
551.4						551.4
552.7						552.7
553.1						553.1
553.7						553.7
554.0						554.0
554.4						554.4
555.2						555.2
555.6						555.6
556.3						556.3
556.9						556.9
557.1						557.1
557.4						557.4
557.8						557.8
558.1						558.1
558.6						558.6
560.0						560.0
561.6						561.6
562.3						562.3
562.9						562.9
563.5						563.5
564.1						564.1
564.8						564.8
565.4						565.4
566.0						566.0
566.7						566.7
567.1						567.1
568.6						568.6
569.9						569.9
570.5						570.5
571.1						571.1
571.7						571.7
572.4						572.4
573.0						573.0
574.3						574.3
574.9						574.9
575.5						575.5
576.2						576.2
576.8						576.8
577.4						577.4
578.1						578.1
579.3						579.3
580.6						580.6
581.9						581.9
582.5						582.5

R/V ACANIA

07 AUG / 09 AUG 79

CHEMICAL MESOSCALE(CRUISE VIII)

TIME GMT	LATITUDE NORTH	LONGITUDE WEST	DISTANCE KM	NO3 UM	PO4 UM	NUTR-RATIO NO3/PO4	ATP NG/L	CHL A MG/M3	TEMP C
0900	35 49.2	122 46.6	583.1						15.4
			583.8						15.4
			584.4						15.4
			585.0						15.4
			585.7						15.4
			586.3						15.4
			586.9						15.4
			587.6						15.4
			588.2						15.4
			588.8						15.4
			589.5						15.4
			590.1						15.4
1000	35 56.5	122 38.9	591.4						15.4
			592.0						15.4
			592.6						15.4
			593.2						15.4
			593.8						15.4
			594.4						15.4
			595.0						15.4
			595.6						15.4
			596.2						15.4
			597.1						15.4
			598.7						15.4
			599.0						15.4
			599.6						15.4
			600.2						15.4
			600.9						15.4
			601.5						15.4
			602.1						15.4
			602.8						15.4
			603.4						15.4
			604.0						15.4
			604.7						15.4
			605.3						15.4
			606.0						15.4
			606.6						15.4
			607.2						15.4
			607.9						15.4
			608.5						15.4
			609.1						15.4
			609.8						15.4
			610.4						15.4
			611.0						15.4
			611.7						15.4
			612.3						15.4
			612.9						15.4
			613.6						15.4
			614.2						15.4



# CHEMICAL MESOSCALE CRUISE VIIII

CHL A  
MG/M3

TEAP  
D=7.3  
14.3

1100	36	3.1	122	32.2
------	----	-----	-----	------

TC/PC	CHL A MG/M3	ATP NG/L	MUTR.RAT ID NO3/PU+	PO4 UM	NO3 UM	DISTANCE KM
0	61.4	0	0	0	0	61.4
1	61.5	0	0	0	0	61.5
2	61.6	0	0	0	0	61.6
3	61.7	0	0	0	0	61.7
4	61.8	0	0	0	0	61.8
5	61.9	0	0	0	0	61.9
6	62.0	0	0	0	0	62.0
7	62.1	0	0	0	0	62.1
8	62.2	0	0	0	0	62.2
9	62.3	0	0	0	0	62.3
10	62.4	0	0	0	0	62.4
11	62.5	0	0	0	0	62.5
12	62.6	0	0	0	0	62.6
13	62.7	0	0	0	0	62.7
14	62.8	0	0	0	0	62.8
15	62.9	0	0	0	0	62.9
16	63.0	0	0	0	0	63.0
17	63.1	0	0	0	0	63.1
18	63.2	0	0	0	0	63.2
19	63.3	0	0	0	0	63.3
20	63.4	0	0	0	0	63.4
21	63.5	0	0	0	0	63.5
22	63.6	0	0	0	0	63.6
23	63.7	0	0	0	0	63.7
24	63.8	0	0	0	0	63.8
25	63.9	0	0	0	0	63.9
26	64.0	0	0	0	0	64.0
27	64.1	0	0	0	0	64.1
28	64.2	0	0	0	0	64.2
29	64.3	0	0	0	0	64.3
30	64.4	0	0	0	0	64.4
31	64.5	0	0	0	0	64.5
32	64.6	0	0	0	0	64.6
33	64.7	0	0	0	0	64.7
34	64.8	0	0	0	0	64.8
35	64.9	0	0	0	0	64.9
36	65.0	0	0	0	0	65.0
37	65.1	0	0	0	0	65.1
38	65.2	0	0	0	0	65.2
39	65.3	0	0	0	0	65.3
40	65.4	0	0	0	0	65.4
41	65.5	0	0	0	0	65.5
42	65.6	0	0	0	0	65.6
43	65.7	0	0	0	0	65.7
44	65.8	0	0	0	0	65.8
45	65.9	0	0	0	0	65.9
46	66.0	0	0	0	0	66.0
47	66.1	0	0	0	0	66.1
48	66.2	0	0	0	0	66.2
49	66.3	0	0	0	0	66.3
50	66.4	0	0	0	0	66.4
51	66.5	0	0	0	0	66.5
52	66.6	0	0	0	0	66.6
53	66.7	0	0	0	0	66.7
54	66.8	0	0	0	0	66.8
55	66.9	0	0	0	0	66.9
56	67.0	0	0	0	0	67.0
57	67.1	0	0	0	0	67.1
58	67.2	0	0	0	0	67.2
59	67.3	0	0	0	0	67.3
60	67.4	0	0	0	0	67.4
61	67.5	0	0	0	0	67.5
62	67.6	0	0	0	0	67.6
63	67.7	0	0	0	0	67.7
64	67.8	0	0	0	0	67.8
65	67.9	0	0	0	0	67.9
66	68.0	0	0	0	0	68.0
67	68.1	0	0	0	0	68.1
68	68.2	0	0	0	0	68.2
69	68.3	0	0	0	0	68.3
70	68.4	0	0	0	0	68.4
71	68.5	0	0	0	0	68.5
72	68.6	0	0	0	0	68.6
73	68.7	0	0	0	0	68.7


# CHEMICAL MESOSCAL CRUISE VILLI


CHL  
PC/03

1300	36 16.7	122 17.6
------	---------	----------


DISTANCE KI	N03 UM	PO4 UM	MUTR.RATIO N03/PO4	ATP MG/L	CHL A MG/100
646.5					
647.1					
648.4					
649.0					
649.7					
650.9					
651.6					
652.2					
652.9					
653.5					
654.1					
654.8					
655.4					
656.0					
656.7					
657.3					
658.0					
659.2					
659.8					
660.5					
661.1					
661.7					
662.4					
663.0					
663.6					
664.3					
664.9					
665.5					
666.2					
666.8					
667.4					
668.1					
668.7					
669.3					
670.0					
670.6					
671.2					
671.9					
672.5					
673.1					
673.8					
674.4					
675.0					
675.7					
676.3					
676.9					
677.6					


# APPENDIX B







31 MAR 1979  
 1 APR 1979  
 2 APR 1979  
 3 APR 1979  
 4 APR 1979  
 5 APR 1979  
 6 APR 1979  
 7 APR 1979  
 8 APR 1979  
 9 APR 1979  
 10 APR 1979  
 11 APR 1979  
 12 APR 1979  
 13 APR 1979  
 14 APR 1979  
 15 APR 1979  
 16 APR 1979  
 17 APR 1979  
 18 APR 1979  
 19 APR 1979  
 20 APR 1979  
 21 APR 1979  
 22 APR 1979  
 23 APR 1979  
 24 APR 1979  
 25 APR 1979  
 26 APR 1979  
 27 APR 1979  
 28 APR 1979  
 29 APR 1979  
 30 APR 1979  
 1 MAY 1979  
 2 MAY 1979











## BIBLIOGRAPHY

1. Anonymous, Technicon Industrial Systems Technical Publication No. TA 1-0170-20, Operation Manual for the Technicon Auto Analyzer II System, 1972.
2. Anonymous, Technicon Industrial Systems Industrial Method No. 177-72-WM, Ortho Phosphate in Water and Seawater, January 1973.
3. Anonymous, Technicon Industrial Systems Industrial Method No. 175-72-WM, Nitrate and Nitrite in Water and Seawater, January 1973.
4. Anonymous, Technicon Industrial Systems Industrial Method No. 100-70-W/B, Nitrate and Nitrite in Water and Waste Water, January 1978.
5. Bakun, A., NOAA Technical Report NMFS SSRF-671, Coastal Upwelling Indices, West Coast of North America 1946-71, 1973.
6. Banse, K., "The Nitrogen-to-Phosphorus Ratio in the Photic Zone of the Sea and the Elemental Composition of the Plankton," Deep Sea Research, V. 21, pp. 767-771.
7. Barton, E. D., Huyer, A. and Smith, R. L., "Temporal Variations Observed in the Hydrographic Regime Near Cabo Corbeiro in the Northwest African Upwelling Region February to April 1974," Deep Sea Research, V. 24, pp. 7-25, 1977.
- \*8. Broecker, W. S., Chemical Oceanography, Harcourt Brace Jovanovich, pp. 9-13, 1974.
9. Bower, R. L., Gohrband, H. S., Pichel, W. G., Signore, T. L., and Walter, C. C., NOAA Technical Memorandum NESS 78, Satellite Derived Sea Surface Temperatures from NOAA Spacecraft, pp. 1-12 and 35-39, June 1976.
10. Butler, E. I., Knox, S., and Liddicott, M. I., "The Relationship Between Inorganic and Organic Nutrients in Seawater," Journal of the Marine Biological Association of the United Kingdom, V. 59, pp. 239-250, 1979.
11. Cheney, R. E., and Winfrey, D. E., NAVOCEANO Technical Note 3700-56-76, Distribution and Classification of Fronts, p. 206, August 1976.

\*Background literature not cited in the text.

12. Codispoti, L. A., and Friederich, G. E., "Local and Mesoscale Influences on Nutrient Variability in the Northwest African Upwelling Region Near Cabo Corbeiro," Deep Sea Research, V. 25, pp. 751-770, 1977.
13. Codispoti, L. A., Dugdale, R. C., Minas, H. J., "A Comparison of the Nutrient Regimes Off Northwest Africa, Peru and Baja California," CUEA Newsletter, V. 8, pp. 2-25, January 1979.
14. Dugdale, R. C., "Nutrient Limitation in the Sea: Dynamics, Identification, and Significance," Limnology and Oceanography, V. 12, pp. 685-695, 1967.
15. Dugdale, R. C., and Georing, J. J., "Uptake of New and Regenerated Forms of Nitrogen in Primary Productivity," Limnology and Oceanography, V. 12, pp. 196-206, 1962.
16. Fairbridge, R. W., The Encyclopedia of Oceanography, Reinhold Publishing Corporation, 1966.
17. Fett, R. W., La Violetter, P. E., Nestor, M., Nickerson, J. W., Rabe, K., NAVENVPREDRSHFAC Technical Report No. 77-04, Navy Tactical Applications Guide, Vol. II, Environmental Phenomena and Effects, pp. 3B-6, January 1979.
18. Holm-Hansen, O. and Karl, D. M., "Effects of Luciferin Concentration on the Quantitative Assay of ATP Using Crude Luciferase Preparations," Analytical Biochemistry, V. 75, pp. 100-112, 1976.
19. Holm-Hansen, O., "Determination of Microbial Biomass in Ocean Profiles," Limnology and Oceanography, V. 14, pp. 740-747, 1969.
20. Huyer, A., Smith, R. L. and Pillsburg, R. D., "Observations in a Coastal Upwelling Region During a Period of Variable Winds (Oregon Coast, July 1972)," Tethys, V. 6, pp. 1-2, 1974.
21. Huntsman, S. A. and Barber, R. T., "Primary Production Off Northwest Africa the Relationship to Wind and Nutrient Conditions," Deep Sea Research, V. 24, pp. 25-35, 1977.
22. Jones, P. G. W., "The Southern Benguela Current Region in February 1966: Part 1 Chemical Observations with Particular Reference to Upwelling," Deep Sea Research, V. 18, pp. 193-208, 1971.
23. Kelly, J. C., Whitley, T. E. and Dugdale, R. C., "Results of Sea Surface Mapping in the Peru Upwelling System," Limnology and Oceanography, V. 20, pp. 784-794, 1975.

24. Legeckis, R. V., "A Survey of Worldwide Sea Surface Temperature Fronts Detected by Environmental Satellites," Journal of Geophysical Research, V. 83, pp. 4501-4523, 20 September 1978.
25. Lorenzen, C. J., "A Method for the Continuous Measurement of In Vivo Chlorophyll Concentration," Deep Sea Research, V. 13, pp. 223-227, 1966.
26. Paulson, G. P., A Study of Nutrient Variations in the Surface and Mixed Layer of Monterey Bay Using Automatic Analysis Techniques, Master's Thesis, Naval Postgraduate School, Monterey, California, September 1972.
27. Redfield, A. C., "On the Proportions of Organic Derivatives in Seawater and Their Relation to the Composition of Plankton," James Johnstone Memorial Volume, University Press of Liverpool, pp. 176-192, 1934.
28. Redfield, A. C., "The Biological Control of Chemical Factors in the Environment," American Scientist, pp. 205-221, September 1958.
29. Redfield, A. C., Ketchum, B. H., and Richards, F. A., "The Influence of Organisms on the Composition of Seawater," in N. M. Hill (ed.), The Sea, V. 2, pp. 26-27, Interscience Publishers, 1963.
30. Reid, J. L., Roden, G. I., and Wyllie, J. G., "Studies of the California Currents Systems," California Cooperative Oceanic Fisheries Investigation, Progress Report, pp. 27-56, 1 July 1956-1 January 1958.
31. Richards, F. A., "Anoxic Basins and Fjords," in J. P. Riley and G. Skirrow (ed.), Chemical Oceanography, V. 1, Academic Press, pp. 611-645, 1965.
32. Riley, J. P., and Skirrow, G., Chemical Oceanography, 2nd Ed., V. 3, Academic Press, 1975.
33. SCOR Working Group 36 (with ACMRR and ACOMR), Coastal Upwelling Processes, Report of 2nd Meeting, Kiel, FRG 24-28 June 1974, CUEA Newsletter, V. 4, pp. 12-20, September 1975.
34. Shaffer, G., "On Quasi-Steady Three-Dimensional Coastal Upwelling Circulation," CUEA Newsletter, V. 5, pp. 25-29, March 1976.
- \*35. Sherman, J. W., III, "Current and Future Satellites for Ocean Monitoring," In: Proceedings of the Eleventh International Symposium on Remote Sensing of Environment, Environmental Research Institute of Michigan, Ann Arbor, Michigan, V. 1, pp. 288-299, 1977.

36. Smith, S. L., and Whitley, T. E., "The Role of Zooplankton in the Regeneration of Nitrogen in a Coastal Upwelling System Off Northwest Africa," Deep Sea Research, V. 24, pp. 49-56, 1977.
37. Strickland, J. D. H. and Parsons, T. R., A Practical Handbook of Seawater Analysis, J. C. Stevenson (Ed.), Queen's Printer and Comptroller of Stationery, pp. 49-62 and 71-76, 1968.
38. Sverdrup, H. U., Johnson, M. W., and Fleming, R. H., The Oceans, Their Physics, Chemistry, and General Biology, Prentice Hall, 1942.
39. Thomas, W. H., and Siebert, D. L. R., "Distribution of Nitrate, Nitrite, Phosphate, and Silicate in the California Current Region, 1969," California Cooperative Oceanic Fisheries Investigations, V. 20, June 1974.
40. Thomas, W. H., "Phytoplankton Nutrient Enrichment Experiments Off Baja California and in the Eastern Equatorial Pacific Ocean," Journal of the Fisheries Research Board of Canada, V. 26, pp. 1133-1145, 1969.
41. Traganza, E. D., Chemical Mesoscale and ATP-Biomass Correlated with Satellite Imagery of Oceanic Fronts, Interim Report on Proposal Submitted to Office of Naval Research, Code 482, 27 June 1978.
42. Traganza, E. D., Naval Postgraduate School Technical Report No. 68-79-008, "The Use of Temperature and Color in Satellite Detection of Ocean Fronts and Eddies for ASW Applications," February 1979.
43. Traganza, E. D., Nestor, D. A., and McDonald, A. K., "Satellite Observations of a Nutrient Upwelling Off the Coast of California," (in press Journal of Geophysical Research), 1980.
44. Treguer, P., and LeCorre, P., "The Ratios of Nitrate Phosphate, and Silicate During Uptake and Regeneration Phases of the Moroccan Upwelling Regime," Deep Sea Research, V. 23A, pp. 163-184, 1977.
- \*45. Wiebe, P. H., "Gulf Stream Cold Core Rings: Large Scale Interaction Sites for Open Ocean Plankton Communities," Deep Sea Research, V. 23, pp. 695-710, 1978.

# INITIAL DISTRIBUTION LIST

	No. Copies
1. Defense Technical Information Center Cameron Station Alexandria, Virginia 22314	2
2. Library, Code 0142 Naval Postgraduate School Monterey, California 93940	2
3. Chairman, Code 68 Department of Oceanography Naval Postgraduate School Monterey, California 93940	3
4. Director Naval Oceanography Division (OP952) Navy Department Washington, DC 20350	1
5. Office of Naval Research Code 480 Naval Ocean Research and Development Activity NSTL Station, Mississippi 39529	1
6. Dr. Robert E. Stevenson Scientific Liaison Office, ONR Scripps Institution of Oceanography La Jolla, California 92037	1
7. SIO Library University of California, San Diego P.O. Box 2367 La Jolla, California 92037	1
8. Department of Oceanography Library University of Washington Seattle, Washington 98105	1
9. Department of Oceanography Library Oregon State University Corvallis, Oregon 97331	1
10. Commanding Officer Fleet Numerical Weather Central Monterey, California 93940	1

11. Commanding Officer 1  
Naval Environmental Prediction Research  
Facility  
Monterey, California 93940
12. Commander 1  
Oceanographic Systems Pacific  
Box 1390  
Pearl Harbor, Hawaii 96860
13. Commanding Officer 1  
Naval Ocean Research and Development  
Activity  
Dr. Edward Green  
Code 482  
NSTL Station, Mississippi 39529
14. Commander 1  
Naval Oceanography Command  
NSTL Station, Mississippi 39529
15. Commanding Officer 1  
Naval Oceanographic Office  
NSTL Station, Mississippi 39529
16. Dr. E. D. Traganza, Code 68Tg 5  
Department of Oceanography  
Naval Postgraduate School  
Monterey, California 93940
17. Mr. Jerry Norton, Code 68 1  
Department of Oceanography  
Naval Postgraduate School  
Monterey, California 93940
18. LT Don A. Nestor 1  
Helicopter Antisubmarine Squadron One  
Naval Air Station  
Jacksonville, Florida 32212
19. Mr. Laurence Breaker 1  
National Environmental Satellite Service  
660 Price Avenue  
Redwood City, California 94063
20. Mr. Roland Nagle 1  
Naval Environmental Prediction Research  
Facility  
Monterey, California 93940
21. Mr. Morton Kronengold 1  
Institute for Acoustical Research  
615 SW 2nd Avenue  
Miami, Florida 33130

- |     |  |   |
|-----|--|---|
| 22. | Commander<br>Area ASW Forces Sixth Fleet<br>LCDR John W. Conrad<br>U.S. Naval Support Activity, Naples<br>FPO New York 09521 | 3 |
| 23. | Mr. Rolf J. Berger, Code 68<br>Department of Oceanography<br>Naval Postgraduate School<br>Monterey, California 93940         | 1 |
| 24. | LT Walter Hanson, Code 35<br>Department of Oceanography<br>Naval Postgraduate School<br>Monterey, California 93940           | 1 |
| 25. | Ms. Andrea McDonald, Code 61<br>Department of Chemistry<br>Naval Postgraduate School<br>Monterey, California 93940           | 1 |
| 26. | Captain W. W. Reynolds, Code 35<br>Department of Oceanography;<br>Naval Postgraduate School<br>Monterey, California 93940    | 1 |
| 27. | Dr. Stevens P. Tucker, Code 68Tx<br>Department of Oceanography<br>Naval Postgraduate School<br>Monterey, California 93940    | 1 |